

**DUSP2 Antibody (C-term) Blocking Peptide**  
Synthetic peptide  
Catalog # BP9084b**Specification****DUSP2 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [Q05923](#)**DUSP2 Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 1844

**Other Names**

Dual specificity protein phosphatase 2, Dual specificity protein phosphatase PAC-1, DUSP2, PAC1

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP9084b](/products/AP9084b) was selected from the C-term region of human DUSP2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**DUSP2 Antibody (C-term) Blocking Peptide - Protein Information**

Name DUSP2

**DUSP2 Antibody (C-term) Blocking Peptide - Background**

DUSP2 is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli.

**DUSP2 Antibody (C-term) Blocking Peptide - References**

Caunt,C.J., et.al., J. Biol. Chem. 283 (10), 6241-6252 (2008)Baranyai,R., et.al., Neuropsychobiology 57 (3), 146-150 (2008)

**Synonyms PAC1****Function**

Regulates mitogenic signal transduction by dephosphorylating both Thr and Tyr residues on MAP kinases ERK1 and ERK2.

**Cellular Location**

Nucleus.

**Tissue Location**

Expressed in hematopoietic tissues.

**DUSP2 Antibody (C-term) Blocking Peptide  
- Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)