


## Phospho-ADAM 17 (Thr735) Ab

Cat.#: AF3361  
Size: 100ul,200ul


Concn.: 1mg/ml  
Source: Rabbit

Mol.Wt.: 93kDa  
Clonality: Polyclonal

|                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|-------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application:                  | WB 1:500-1:2000, IF/ICC 1:100-1:500                                                                                                                                                                                                                                                                                                                                                                                                            |
| Reactivity:                   | Human,Mouse,Rat                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Purification:                 | The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.                                                                                                                                                                                                                                                                                              |
| Specificity:                  | Phospho-ADAM 17 (Thr735) Ab detects endogenous levels of ADAM 17 only when phosphorylated at Threonine 735.                                                                                                                                                                                                                                                                                                                                    |
| Immunogen:                    | A synthesized peptide derived from human ADAM 17 around the phosphorylation site of Threonine 735.                                                                                                                                                                                                                                                                                                                                             |
| Uniprot:                      | P78536                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Description:                  | This gene encodes a disintegrin and metalloprotease (ADAM) domain 17, which is a member of the ADAM protein family. Members of this family are membrane-anchored proteins structurally related to snake venom disintegrins, and have been implicated in a variety of biologic processes involving cell-cell and cell-matrix interactions, including fertilization, muscle development, and neurogenesis.                                       |
| Subcellular Location:         | Membrane.                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Tissue Specificity:           | Ubiquitously expressed. Expressed at highest levels in adult heart, placenta, skeletal muscle, pancreas, spleen, thymus, prostate, testes, ovary and small intestine, and in fetal brain, lung, liver and kidney.                                                                                                                                                                                                                              |
| Similarity:                   | Must be membrane anchored to cleave the different substrates. The cytoplasmic domain is not required for the this activity. Only the catalytic domain is essential to shed TNF and p75 TNFR (By similarity).The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme. |
| Storage Condition and Buffer: | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.                                                                                                                                                                                                                                                                                   |



Western blot analysis of Phospho-ADAM 17 (Thr735) expression in various lysates



Western blot analysis of ADAM 17 phosphorylation expression in K562UV treated whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



AF3361 staining HeLa cells by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab (Cat.# S0006), diluted at 1/600, was used as secondary Ab.

**IMPORTANT:** For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.