

## Phospho-NF kappaB p100/p52 (Ser869) Ab

Cat.#: AF3374	Concn.: 1mg/ml	Mol.Wt.: 100kDa
Size: 100ul,200ul	Source: Rabbit	Clonality: Polyclonal

**Application:** WB 1:500-1:2000 IHC 1:50-1:200 IP, IF/ICC 1:100-1:500

**Reactivity:** Human,Mouse,Rat

**Purification:** The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

**Specificity:** Phospho-NF- kappaB p100/p52 (Ser869) Ab detects endogenous levels of NF- kappaB p100/p52 only when phosphorylated at Serine 869.

**Immunogen:** A synthesized peptide derived from human NF- kappaB p100/p52 around the phosphorylation site of Serine 869.


**Uniprot:** Q00653

**Description:** NFkB-p100 a transcription factor of the nuclear factor-kappaB ( NFkB) group. Precursor of the p52 subunit of the nuclear factor NF-kappa-B, which binds to the kappa-B consensus sequence 5'-GGRNNYCC-3', located in the enhancer region of genes involved in immune response and acute phase reactions.


**Subcellular Location:** Cytoplasmic and Nuclear

**Similarity:** The C-terminus of p100 might be involved in cytoplasmic retention, inhibition of DNA-binding by p52 homodimers, and/or transcription activation.The glycine-rich region (GRR) appears to be a critical element in the generation of p52.


**Storage Condition and Buffer:** Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of Phospho-NF kappaB p100/p52 (Ser869) expression in various lysates



Western blot analysis of NF- kappaB p100/p52 phosphorylation expression in TNF- $\alpha$  treated MDA-MB-435 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



AF3374 at 1/200 staining human Fallopian tube tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.



AF3374 staining RAW264.7 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

**IMPORTANT:** For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.