


Phospho-PKCD (Tyr64) Ab

Cat.#: AF3410
 Size: 100ul,200ul


Concn.: 1mg/ml
 Source: Rabbit

Mol.Wt.: 78kDa
 Clonality: Polyclonal

| | |
|-------------------------------|---|
| Application: | WB 1:500-1:2000, IF/ICC 1:100-1:500 |
| Reactivity: | Human,Mouse,Rat |
| Purification: | The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns. |
| Specificity: | Phospho-PKCD (Tyr64) Ab detects endogenous levels of PKCD only when phosphorylated at Tyrosine 64. |
| Immunogen: | A synthesized peptide derived from human PKCD around the phosphorylation site of Tyrosine 64. |
| Uniprot: | Q05655 |
| Description: | Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. |
| Subcellular Location: | Cytoplasm. Membrane. |
| Similarity: | The C1 domain, containing the phorbol ester/DAG-type region 1 (C1A) and 2 (C1B), is the diacylglycerol sensor.The C2 domain is a non-calcium binding domain. It binds to proteins containing phosphotyrosine in a sequence-specific manner.Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily. |
| Storage Condition and Buffer: | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt. |



Western blot analysis of Phospho-PKCD (Tyr64) expression in various lysates



Western blot analysis of PKCD phosphorylation expression in PMA treated HepG2 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



AF3410 staining NIH-3T3 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.