

Phospho-IL-10R alpha (Tyr496) Ab

Cat.#: AF3435	Concn.: 1mg/ml	Mol.Wt.: 63kDa
Size: 100ul,200ul	Source: Rabbit	Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500

Reactivity: Human

Purification: The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

Specificity: Phospho-IL-10R alpha (Tyr496) Ab detects endogenous levels of IL-10R alpha only when phosphorylated at Tyrosine 496.

Immunogen: A synthesized peptide derived from human IL-10R alpha around the phosphorylation site of Tyrosine 496.

Uniprot: Q13651


Description: The protein encoded by this gene is a receptor for interleukin 10. This protein is structurally related to interferon receptors. It has been shown to mediate the immunosuppressive signal of interleukin 10, and thus inhibits the synthesis of proinflammatory cytokines.

Subcellular Location: Membrane.


Tissue Specificity: Spleen, thymus, and PBMC. Weak expression in pancreas, skeletal muscle, brain, heart, and kidney. Placenta, lung, and liver showed intermediate levels. Monocytes, B-cells, large granular lymphocytes, and T-cells express high levels.

Similarity: Belongs to the type II cytokine receptor family.


Storage Condition and Buffer: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of Phospho-IL-10R alpha (Tyr496) Ab expression in HuvEc cells lysates.The lane on the right is treated with the antigen-specific peptide.



Western blot analysis of IL-10R alpha phosphorylation expression in HuvEc whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



AF3435 staining HuvEc by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.