


Phospho-STAT4 (Tyr693) Ab

Cat.#: AF3441
 Size: 100ul,200ul


Concn.: 1mg/ml
 Source: Rabbit

Mol.Wt.: 86kDa
 Clonality: Polyclonal


- Application:** WB 1:500-1:2000 IHC 1:50-1:200 IP, IF/ICC 1:100-1:500
- Reactivity:** Human,Mouse,Rat
- Purification:** The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
- Specificity:** Phospho-STAT4 (Tyr693) Ab detects endogenous levels of STAT4 only when phosphorylated at Tyrosine 693.
- Immunogen:** A synthesized peptide derived from human STAT4 around the phosphorylation site of Tyrosine 693.
- Uniprot:** Q14765
- Description:** The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators.
- Subcellular Location:** Cytoplasm. Nucleus. Translocated into the nucleus in response to phosphorylation.
- Similarity:** Belongs to the transcription factor STAT family.
- Storage Condition and Buffer:** Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of Phospho-STAT4 (Tyr693) expression in Mouse heart tissue lysate



Western blot analysis of STAT4 phosphorylation expression in IL-4 treated HeLa whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



AF3441 at 1/200 staining human kidney tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.



AF3441 staining HeLa by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.