

## PKM1/2 Ab

Cat.#: AF4031 Concn.: 1mg/ml Mol.Wt.: 60kd Size: 50ul,100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000

Reactivity: Human

Purification: The antiserum was purified by peptide affinity

chromatography using SulfoLink™ Coupling Resin (Thermo

Fisher Scientific).

Specificity: The Ab detects endogenous levels of total PKM1/2 protein.

Immunogen: Peptide sequence around aa. 36~40(D-S-P-P-I) derived from

Human PKM1/2.

Uniprot: P14618

Description: Glycolytic enzyme that catalyzes the transfer of a

phosphoryl group from phosphoenolpyruvate (PEP) to ADP, generating ATP. Stimulates POU5F1-mediated transcriptional activation. Plays a general role in caspase independent cell death of tumor cells. The ratio between the highly active tetrameric form and nearly inactive dimeric form determines whether glucose carbons are channeled to biosynthetic processes or used for glycolytic ATP production. The transition between the 2 forms contributes to the control of glycolysis and is important for tumor cell proliferation and

survival.

Subcellular Location: Cytoplasm. Nucleus. Translocates to the nucleus in response

to different apoptotic stimuli. Nuclear translocation is sufficient to induce cell death that is caspase independent, isoform-specific and independent of its enzymatic activity.

Tissue Specificity: Specifically expressed in proliferating cells, such as

embryonic stem cells, embryonic carcinoma cells, as well as

cancer cells.

Similarity: Belongs to the pyruvate kinase family.

Storage Condition and

Buffer:


Supplied at 1.0mg/mL in phosphate buffered saline (without

Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide

and 50% glycerol.



## Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



Western blot analysis of extracts from HepG2 and 293 cells using PKM1/2

IMPORTANT: For western blot, incubate membrane with diluted primary Ab
in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking,
overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.