



Goat anti-TAF8 / TAFII43 Antibody

Item Number	dAP-1986
Target Molecule	Principle Name: TAF8 / TAFII43; Official Symbol: TAF8; All Names and Symbols: TAF8; TAF8 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 43kDa; 43; FLJ32821; II; TAF; TAFII43; TBN; OTTHUMP00000016392; TAF(II)43; TAF8 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 50 kD; TATA box binding pr; Accession Number (s): NP_612639.2; Human Gene ID(s): 129685; Non-Human GeneID(s): 63856 (mouse) 316216 (rat)
Immunogen	KSYCEHTARTQPT, is from internal region
Applications	Pep ELISA Species Tested:
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 16000.
Western Blot	Western Blot: Preliminary experiments gave an approx 38kDa band in Human Brain (Amygdala, Cerebellum, Hippocampus) lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe
IHC	
Reference	Reference(s): Guermah M, Ge K, Chiang CM, Roeder RG, The TBN protein, which is essential for early embryonic mouse development, is an inducible TAFII implicated in adipogenesis. Molecular cell 2003 Oct 12 (4): 991-1001..PMID: 14580349->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**