



## Goat anti-Npc1(mouse) Antibody

<b>Item Number</b>	dAP-2150
<b>Target Molecule</b>	Principle Name: Npc1(mouse); Official Symbol: Npc1; All Names and Symbols: Npc1; Niemann Pick type C1; A430089E03Rik; C85354; D18ErtD139e; D18ErtD723e; lcsd; nmf164; spm; niemann-Pick C1 protein; sphingomyelinosis; Accession Number (s): NP_032746.2; Human Gene ID(s): ; Non-Human GeneID(s): 18145 (mouse)
<b>Immunogen</b>	TYHTILKTSADYTD, is from internal region
<b>Applications</b>	Pep ELISA Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 16000.
<b>Western Blot</b>	Western Blot: Preliminary experiments gave an approx 36kDa band in lysates of cell lines NIH3T3 and NSO after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculate
<b>IHC</b>	
<b>Reference</b>	Reference(s): Tang Y, Leao IC, Coleman EM, Broughton RS, Hildreth JE, Deficiency of niemann-pick type C-1 protein impairs release of human immunodeficiency virus type 1 and results in Gag accumulation in late endosomal/lysosomal compartments. Journal of virology 2009 Aug 83 (16): 7982-95..PMID: 19474101-

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**