



Goat anti-contactin 1 Antibody

Item Number dAP-2999

Principle Name: contactin 1; Official Symbol: CNTN1; All Names and Symbols: CNTN1; contactin 1; F3; **Target Molecule**

GP135; contactin-1; glycoprotein gP135; neural cell surface protein F3; Accession Number (s): NP_001834.2; NP_778203.1; NP_001242992.1; Human Gene ID(s): 1272; Non-Human GeneID(s): 12805

(mouse) 117258 (rat)

Immunogen ENIRGKDKHQAR, is from internal region

This antibody is expected to recognize all reported isoforms (NP 001834.2; NP 778203.1;

NP_001242992.1). Reported variants represent identical protein: NP_001242993.1, NP_001242992.1.

Applications

Species Tested: Human, Mouse, Rat, Pig

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

using the immunizing peptide.

Supplied As lyophilized powder of 50ug or 100ug IgG; Reconsititute IgG with 100ul or 200ul sterile DI Water and final

product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

Aliquot and store at -20°C. Minimize freezing and thawing.

Peptide ELISA: antibody detection limit dilution 1 to 32000. Peptide ELISA

Western Blot Western Blot: Approx 140kDa band observed in Human, Mouse, Rat and Pig Brain lysates (calculated MW

of 113kDa according to NP 001834.2). The observed molecular weight corresponds to the glycosylated

form. Recommended concentration: 0.3-1µg/ml.

IHC

Reference(s): Lamprianou S, Chatzopoulou E, Thomas JL, Bouyain S, Harroch S. A complex between Reference

contactin-1 and the protein tyrosine phosphatase PTPRZ controls the development of oligodendrocyte

precursor cells. Proc Natl Acad Sci U S A. 2011 Oct 18;108(42):17498-503..PMID: 21969550->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only