



Mouse Monoclonal Antibody to CD207

Cataloge Number sAP-1668

Target Molecule Name: CD207

Aliases: CLEC4K

MW: 36.7kDa

Entrez Gene ID: 50489

Descrption The protein encoded by this gene is expressed only in Langerhans cells which are immature dendritic cells

of the epidermis and mucosa. It is localized in the Birbeck granules, organelles present in the cytoplasm of Langerhans cells and consisting of superimposed and zippered membranes. It is a C-type lectin with mannose binding specificity, and it has been proposed that mannose binding by this protein leads to internalization of antigen into Birbeck granules and providing access to a nonclassical antigen-processing pathway.

Mutations in this gene result in Birbeck granules deficiency or loss of sugar binding activity.

Immunogen Purified recombinant fragment of human CD207 (AA: extra 155-328) expressed in E. Coli.

Recitative Species Human;

Clone MM1C5A1

Size and Concentration 100µg/1mg/ml

Supplied as Lyophilized Powder from 100µl of Purified antibody in PBS with 0.05% sodium azide

Reconstitution/Storages Reconstituted with 100µl sterile DI H2O, at stored at 4°C or -20°C for short or long term storage

Applications ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; ICC: N to A; FCM: 1 to 200 - 1 to 400; IHC: N to A

Shipping Regular FEDEX overnight shipment (ambient temperature)

Reference 1.Immunol Cell Biol. 2015 Oct;93(9):815-24.2.J Immunol. 2011 Feb 1;186(3):1377-83.

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only