# pMXs-IRES-Puro Retroviral Vector

CATALOG NUMBER: RTV-014 STORAGE: -20°C

# **QUANTITY AND CONCENTRATION:** 10 $\mu$ g at 0.25 $\mu$ g/ $\mu$ L in TE

#### **Background**

Retroviruses are efficient tools for delivering heritable genes into the genome of dividing cells. Cell Biolabs' pMXs-IRES-Puro retroviral vector (also known as pMXs-IP) is based on Moloney murine leukemia virus (MMLV). The vector provides the viral package signal, transcription and processing elements, and MCS for cloning of a target gene. The viral *env* gene, produced by the package cell line, encodes the envelope protein, which determines the viral infectivity range. Transfection into a package cell line produces high-titer, replication-incompetent viruses. In addition to transfer and expression of exogenous genes in mammalian cells, recently, retroviruses have been used to express silencing RNAs (siRNA) to decrease the expression of target genes both *in vitro* and *in vivo*.

The vector contains the ampicillin-resistance gene, MMLV LTRs, package signal and MCS for cloning of your gene of interest (Figure 1).




Figure 1. Schematic representation of pMXs-IRES-Puro retroviral vector.

MCS:

- Enzyme Sites: 5'-BamHI, EcoRI, XhoI, NotI, SnaBI-3'
- MCS Sequence: TTAATTAA<u>GGATCC</u>CAGTGTGGTGGTACGG<u>GAATTC</u>CTGCAGGC<u>CTCGAG</u>GGCCGGC GCGCC<u>GCGGCCGCTACGTA</u>AATT---IRES---puro---



#### **Safety Consideration**

Remember that you will be working with samples containing infectious virus. Follow the recommended NIH guidelines for all materials containing BSL-2 organisms. Always wear gloves, use filtered tips and work under a biosafety hood.

#### **References**

1. Kitamura T., et al., (2003) Exp. Hematol. 31, 1007-1014.

## **Recent Product Citations**

- 1. Koso, H. et al. (2014). Identification of FoxR2 as an oncogene in medulloblastoma. *Cancer Res.* **74**:2351-2361.
- Sugatani, T. et al. (2011). A microRNA Expression Signature of Osteoclastogenesis. *Blood*. 117:3648-3657.
- 3. Sugatani, T. and K. Hruska (2009). Impaired microRNA pathways diminish osteoclast differentiation and function. *J. Biol. Chem.* **284**:4667-4678.

## **License Information**

This product is licensed from the University of Tokyo.

#### **Warranty**

These products are warranted to perform as described in their labeling and in Cell Biolabs literature when used in accordance with their instructions. THERE ARE NO WARRANTIES THAT EXTEND BEYOND THIS EXPRESSED WARRANTY AND CELL BIOLABS DISCLAIMS ANY IMPLIED WARRANTY OF MERCHANTABILITY OR WARRANTY OF FITNESS FOR PARTICULAR PURPOSE. CELL BIOLABS's sole obligation and purchaser's exclusive remedy for breach of this warranty shall be, at the option of CELL BIOLABS, to repair or replace the products. In no event shall CELL BIOLABS be liable for any proximate, incidental or consequential damages in connection with the products.

#### This product is for RESEARCH USE ONLY; not for use in diagnostic procedures.

# **Contact Information**

Cell Biolabs, Inc. 7758 Arjons Drive San Diego, CA 92126 Worldwide: +1 858-271-6500 USA Toll-Free: 1-888-CBL-0505 E-mail: <u>tech@cellbiolabs.com</u> www.cellbiolabs.com

©2008-2014: Cell Biolabs, Inc. - All rights reserved. No part of these works may be reproduced in any form without permissions in writing.

