

## PLA2G2A

## Recombinant Human Secreted Phospholipase A2 group IIA His

**Catalog No.** CSI12691 **Quantity**: 2 μg

CSI12692 10 μg CSI12693 1.0 mg

Alternate Names: MOM1, PLA2, PLA2B, PLA2L, PLA2S, PLAS1, sPLA2, Phospholipase A2 membrane

associated, EC 3.1.1.4, Phosphatidylcholine 2-acylhydrolase, Group IIA phospholipase A2, GIIC sPLA2, Non-pancreatic secretory phospholipase A2, NPS-PLA2, sPLA2-IIA,

PLA2G2A.

**Description:** Phospholipase A2 (PLA2) catalyzes the hydrolysis of the sn-2 position of membrane

glycerophospholipids to liberate arachidonic acid (AA), a precursor of eicosanoids including prostaglandins and leukotrienes. The same reaction also produces

lysophosholipids, which represent another class of lipid mediators.

The secretory PLA2 (sPLA2) family, in which 10 isozymes have been identified, consists of low molecular weight, Ca2+-requiring secretory enzymes that have been implicated in

a number of biological processes, such as modification of eicosanoid generation,

inflammation, and host defense.

This enzyme has been proposed to hydrolyze phosphatidylcholine (PC) in lipoproteins to

liberate lyso- PC and free fatty acids in the arterial wall, thereby facilitating the accumulation of bioactive lipids and modified lipoproteins in atherosclerotic foci.

In mice, sPLA2 expression significantly influences HDL particle size and composition and demonstrate that an induction of sPLA2 is required for the decrease in plasma HDL cholesterol in response to inflammatory stimuli. Instillation of bacteria into the bronchi was associated with surfactant degradation and a decrease in large:small ratio of

surfactant aggregates in rats.

sPLA2-IIA can exert beneficial action in the context of infectious diseases since recent studies have shown that this enzyme exhibits potent bactericidal effects. Induction of the synthesis of sPLA2-IIA is generally initiated by endotoxin and a limited number of

cytokines via paracrine and/or autocrine processes.

Description Secreted Phospholipase A2-IIA Human Recombinant is manufactured with N-terminal fusion of HisTag. PLA2G2A His-Tagged Fusion Protein is 15.8 kDa containing 124 amino acid residues of the human secreted phospholipase A2-IIA and 16 additional

amino acid residues - HisTag (underlined).

MRGSHHHHHH GMASHMNLVN FHRMIKLTTG KEAALSYGFY GCHCGVGGRG SPKDATDRCC VTHDCCYKRL EKRGCGTKFL SYKFSNSGSR ITCAKQDSCR

SQLCECDKAA ATCFARNKTT YNKKYQYYSN KHCRGSTPRC

 Gene ID:
 5320

 Source:
 E. coli

Molecular Weight: PLA2G2A His-Tagged Fusion Protein is 15.8 kDa containing 124 amino acid residues of

Toll Free: 888-769-1246

Phone: 781-828-0610


Fax: 781-828-0542

the Human Secreted Phospholipase A2-IIA and 16 additional amino acid residues.

E-mail: info@cellsciences.com
Website: www.cellsciences.com

**Formulation:** Lyophilized from 0.5 mg/ml in 0.05 M Acetate buffer pH-4.

**Purity:** Greater than 95% as determined by SDS-PAGE. **Physical Appearance:** Sterile Filtered lyophilized (freeze-dried) powder.



## cellsciences.com

Purification: The SETD7 purified by proprietary chromatographic techniques.

Specific Activity: The amino acid sequence of the PLA2G2A is 100% homologous to the amino acid

sequence of the human Secreted Phospholipase A2-IIA.

NLVN FHRMIKLTTG KEAALSYGFY GCHCGVGGRG SPKDATDRCC VTHDCCYKRL Amino Acid Sequence:

EKRGCGTKFL SYKFSNSGSR ITCAKQDSCR SQLCECDKAA ATCFARNKTT

YNKKYQYYSN KHCRGSTPRC

Reconstitution: Add 0.2 ml of 0.1 M Acetate buffer pH-4 and let the lyophilized pellet dissolve completely.

> For conversion into higher pH value, we recommend intensive dilution by relevant buffer to a concentration of 10 µg/ml. In higher concentrations the solubility of this antigen is

limited.

**Purification Method:** Three-step procedure using affinity Ni-NTA chromatography and size exclusion

chromatography before and after refolding.

Store lyophilized protein at -20°C. The lyophilized protein remains stable until the Storage & Stability:

expiration date when stored at -20°C.

Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show

any change after two weeks at 4°C.

Applications: Western blot

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

Toll Free: 888-769-1246 E-mail: info@cellsciences.com Phone: 781-828-0610 Fax: 781-828-0542

Website: www.cellsciences.com