cellsciences.com

Recombinant Hepatitis C Virus NS3 Subtype 1b (aa 1192 -1459) GST

Catalog No.	CSI15766A CSI15766B CSI15766C	Quantity:	100 μg 0.5 mg 1.0 mg
Description:	HCV is a small 50 nm, enveloped, single-stranded, positive sense RNA virus in the family Flaviviridae. HCV has a high rate of replication with approximately one trillion particles produced each day in an infected individual. Due to lack of proofreading by the HCV RNA polymerase, the HCV has an exceptionally high mutation rate, a factor that may help it elude the host's immune response. Hepatitis C virus is classified into six genotypes (1-6) with several subtypes within each genotype. The preponderance and distribution of HCV genotypes varies globally. Genotype is clinically important in determining potential response to interferon-based therapy and the required duration of such therapy. Genotypes (2, 3, 5 and 6). The <i>E.coli</i> derived recombinant protein contains the HCV NS3 immunodominant regions, amino acids 1192-1459. The protein is fused to a GST tag at N-Terminus.		
Source:	E. coli		
Source.	E: 00//		
Formulation:	1.5 M urea + 2 5mM Tris-HC	pH-8, + 0.2% Triton-X + 5	50% Glycerol.
	1.5 M urea + 2 5mM Tris-HC	•	50% Glycerol. hined by 10% PAGE (coomassie
Formulation:	1.5 M urea + 2 5mM Tris-HC HCV NS3 Genotype-1b prote staining).	in is >95% pure as determ	-
Formulation: Purity:	1.5 M urea + 2 5mM Tris-HC HCV NS3 Genotype-1b prote staining).	in is >95% pure as determ in was purified by propriet	nined by 10% PAGE (coomassie
Formulation: Purity: Purification Method:	1.5 M urea + 2 5mM Tris-HC HCV NS3 Genotype-1b prote staining). HCV NS3 Genotype-1b prote	in is >95% pure as determ in was purified by propriet HCV-infected individuals. 4°C for 1 week, should be	nined by 10% PAGE (coomassie ary chromatographic technique.
Formulation: Purity: Purification Method: Specific Activity:	 1.5 M urea + 2 5mM Tris-HC HCV NS3 Genotype-1b protestaining). HCV NS3 Genotype-1b protestain Immunoreactive with sera of HCV-Core although stable at 	in is >95% pure as determ in was purified by propriet HCV-infected individuals. 4°C for 1 week, should be cycles. en is suitable for ELISA ar	ary chromatographic technique. e stored below -18°C.

