

🕜 Tel: +1-301-363-4651 🗵 Email: cusabio@cusabio.com 🙆 Website: www.cusabio.com 🌘

LIAS Antibody

Product CodeCSB-PA012927GA01HUAbbreviationLIASAbbreviationUpon receipt, store at -20°C or -80°C. Avoid repeated freezVinjord No.O43766ImmunogenHuman LIASRaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasIipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)Target NamesLIAS	erol, pH 7.320°C, Avoid freeze / thaw AS;LIP1;MGC23245 ; htibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It hemistry in lipoate biosynthesis.
StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freezUniprot No.O43766ImmunogenHuman LIASRaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	erol, pH 7.320°C, Avoid freeze / thaw AS;LIP1;MGC23245 ; htibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It hemistry in lipoate biosynthesis.
Uniprot No.O43766ImmunogenHuman LIASRaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliasIipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	erol, pH 7.320°C, Avoid freeze / thaw AS;LIP1;MGC23245 ; htibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It hemistry in lipoate biosynthesis.
ImmunogenHuman LIASRaised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	AS;LIP1;MGC23245 ; htibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It hemistry in lipoate biosynthesis.
Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	AS;LIP1;MGC23245 ; ntibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It nemistry in lipoate biosynthesis.
Species ReactivityHuman,Mouse,RatTested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	AS;LIP1;MGC23245 ; htibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It hemistry in lipoate biosynthesis.
Tested ApplicationsELISA,WB,IHCStorage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	AS;LIP1;MGC23245 ; ntibody ic acid synthetases family. It localizes ole in alpha-(+)-lipoic acid synthesis. It nemistry in lipoate biosynthesis.
Storage BufferPBS with 0.1% Sodium Azide, 50% Glycerol, pH 7.320°C cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	AS;LIP1;MGC23245 ; htibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It hemistry in lipoate biosynthesis.
cycles.Purification MethodAntigen Affinity PurifiedIsotypeIgGAliaslipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	AS;LIP1;MGC23245 ; ntibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It nemistry in lipoate biosynthesis.
IsotypeIgGAliaslipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	ntibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It nemistry in lipoate biosynthesis.
Aliaslipoic acid synthetase;LIAS;HUSSY-01;LAS;LIP1;MGC2324Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	ntibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It nemistry in lipoate biosynthesis.
Product TypePurified Rabbit Anti human PolyClonal AntibodySpeciesHomo sapiens (Human)	ntibody ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It nemistry in lipoate biosynthesis.
Species Homo sapiens (Human)	ic acid synthetases family. It localizes role in alpha-(+)-lipoic acid synthesis. It nemistry in lipoate biosynthesis.
	ole in alpha-(+)-lipoic acid synthesis. It nemistry in lipoate biosynthesis.
Target Names LIAS	ole in alpha-(+)-lipoic acid synthesis. It nemistry in lipoate biosynthesis.
	ole in alpha-(+)-lipoic acid synthesis. It nemistry in lipoate biosynthesis.
Target DetailsThis protein belongs to the biotin and lipoic acid synthetases in mitochondrion and plays an important role in alpha-(+)-lip may also function in the sulfur insertion chemistry in lipoate Alternative splicing occurs at this locus and two transcript va distinct isoforms have been identified.	id two transcript variants encoding

1