

Histone H2B Antibody, HRP conjugated

Product CodeCSB-PA355820LB01CSJAbbreviationHistone H2BStorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.P14001ImmunogenRecombinant Crocodylus niloticus Histone H2B protein (2-93AA)Raised InRabbitSpecies ReactivityCairina moschataTested ApplicationsELISARelevanceCore component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require pulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.FormLiquidConjugateHRPStorage BufferSp5%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthersTarget NamesN/A		
StorageUpon receipt, store at -20°C or -80°C. Avoid repeated freeze.Uniprot No.P14001ImmunogenRecombinant Crocodylus niloticus Histone H2B protein (2-93AA)Raised InRabbitSpecies ReactivityCairina moschataTested ApplicationsELISARelevanceCore component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Product Code	CSB-PA355820LB01CSJ
Uniprot No.P14001ImmunogenRecombinant Crocodylus niloticus Histone H2B protein (2-93AA)Raised InRabbitSpecies ReactivityCairina moschataTested ApplicationsELISARelevanceCore component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Abbreviation	Histone H2B
ImmunogenRecombinant Crocodylus niloticus Histone H2B protein (2-93AA)Raised InRabbitSpecies ReactivityCairina moschataTested ApplicationsELISARelevanceCore component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised InRabbitSpecies ReactivityCairina moschataTested ApplicationsELISARelevanceCore component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Uniprot No.	P14001
Species ReactivityCairina moschataEsted ApplicationsELISARelevanceCore component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Immunogen	Recombinant Crocodylus niloticus Histone H2B protein (2-93AA)
Tested ApplicationsELISARelevanceCore component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Reserch AreaOthers	Raised In	Rabbit
RelevanceCore component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Species Reactivity	Cairina moschata
chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.FormLiquidConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Tested Applications	ELISA
ConjugateHRPStorage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Relevance	chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of
Storage BufferPreservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, PH 7.4Purification Method>95%, Protein G purifiedIsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Conjugate	HRP
IsotypeIgGClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Storage Buffer	
ClonalityPolyclonalAliasHistone H2BSpeciesCairina moschata (Muscovy duck)Research AreaOthers	Purification Method	>95%, Protein G purified
Alias Histone H2B Species Cairina moschata (Muscovy duck) Research Area Others	Isotype	lgG
Species Cairina moschata (Muscovy duck) Research Area Others	Clonality	Polyclonal
Research Area Others	Alias	Histone H2B
	Species	Cairina moschata (Muscovy duck)
Target Names N/A	Research Area	Others
	Target Names	N/A