





GPR18 Antibody, HRP conjugated

| Product Code | CSB-PA614525LB01HU |
|----------------------------|--|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q14330 |
| Immunogen | Recombinant Human N-arachidonyl glycine receptor protein (290-331AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA |
| Relevance | Receptor for endocannabinoid N-arachidonyl glycine (NAGIy) (PubMed:16844083, PubMed:24762058, PubMed:27572937). However, conflicting results about the role of NAGIy as an agonist are reported (PubMed:27018161). Can also be activated by plant-derived and synthetic cannabinoid agonists (PubMed:24762058). The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase (PubMed:16844083). May contribute to regulation of the immune system. Is required for normal homeostasis of CD8+ subsets of intraepithelial lymphocytes (IELs) (CD8alphaalpha and CD8alphabeta IELs)in small intstine by supporting preferential migration of CD8alphaalpha T-cells to intraepithelial compartment over lamina propria compartment, and by mediating their reconstitution into small intestine after bone marrow transplant (By similarity). Plays a role in hypotensive responses, mediating reduction in intraocular and blood pressure (By similarity). Mediates NAGly-induced process of reorganization of actin filaments and induction of acrosomal exocytosis (PubMed:27572937). |
| Form | Liquid |
| Conjugate | HRP |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | N-arachidonyl glycine receptor (NAGly receptor) (G-protein coupled receptor 18), GPR18, GPCRW |
| Species | Human |
| Research Area | Neuroscience |
| Target Names | GPR18 |