





## KCTD6 Antibody, HRP conjugated

| Product Code               | CSB-PA822756LB01HU  |
|----------------------------|---|
| Storage                    | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |
| Uniprot No.                | Q8NC69  |
| Immunogen                  | Recombinant Human BTB/POZ domain-containing protein KCTD6 protein (1-237AA)   |
| Raised In                  | Rabbit  |
| Species Reactivity         | Human   |
| <b>Tested Applications</b> | ELISA   |
| Relevance                  | Probable substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex mediating the ubiquitination and subsequent proteasomal degradation of target proteins. Promotes the ubiquitination of HDAC1; the function seems to depend on KCTD11:KCTD6 oligomerization. Can function as antagonist of the Hedgehog pathway by affecting the nuclear transfer of transcription factor GLI1; the function probably occurs via HDAC1 down-regulation, keeping GLI1 acetylated and inactive. Inhibits cell growth and tumorigenicity of medulloblastoma (MDB) (PubMed:21472142). Involved in regulating protein levels of ANK1 isoform Mu17 probably implicating CUL3-dependent proteasomal degradation (PubMed:22573887). |
| Form                       | Liquid  |
| Conjugate                  | HRP   |
| Storage Buffer             | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  |
| <b>Purification Method</b> | >95%, Protein G purified  |
| Isotype                    | IgG   |
| Clonality                  | Polyclonal  |
| Alias                      | BTB/POZ domain-containing protein KCTD6 (KCASH3 protein) (Potassium channel tetramerization domain-containing protein 6), KCTD6   |
| Species                    | Human   |
| Research Area              | Others  |