




# FCRL4 Antibody

|                            |   |
|----------------------------|---|
| <b>Product Code</b>        | CSB-PA836277LA01HU  |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |
| <b>Uniprot No.</b>         | Q96PJ5  |
| <b>Immunogen</b>           | Recombinant Human Fc receptor-like protein 4 protein (20-387AA)   |
| <b>Raised In</b>           | Rabbit  |
| <b>Species Reactivity</b>  | Human   |
| <b>Tested Applications</b> | ELISA, WB; Recommended dilution: WB:1:500-1:2000  |
| <b>Relevance</b>           | May function as an inhibitor of the B-cell receptor signaling. May function in the B-cell-mediated immune response.   |
| <b>Form</b>                | Liquid  |
| <b>Conjugate</b>           | Non-conjugated  |
| <b>Storage Buffer</b>      | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  |
| <b>Purification Method</b> | >95%, Protein G purified  |
| <b>Isotype</b>             | IgG   |
| <b>Clonality</b>           | Polyclonal  |
| <b>Alias</b>               | Fc receptor-like protein 4 (FcR-like protein 4) (FcRL4) (Fc receptor homolog 4) (FcRH4) (IFGP family protein 2) (hIFGP2) (Immune receptor translocation-associated protein 1) (CD antigen CD307d), FCRL4, FCRH4 IFGP2 IRTA1 |
| <b>Species</b>             | Human   |
| <b>Research Area</b>       | Immunology  |
| <b>Target Names</b>        | FCRL4   |

## Image



Western blot  
 All lanes: FCRL4 antibody at 6µg/ml + 293T whole cell lysate  
 Secondary  
 Goat polyclonal to rabbit IgG at 1/10000 dilution  
 Predicted band size: 58, 57 kDa  
 Observed band size: 58 kDa