





## SLC6A15 Antibody, Biotin conjugated

CSB-PA875657LD01HU
Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Q9H2J7
Recombinant Human Sodium-dependent neutral amino acid transporter B(0)AT2 protein (357-452AA)
Rabbit
Human
ELISA
Functions as a sodium-dependent neutral amino acid transporter. Exhibits preference for the branched-chain amino acids, particularly leucine, valine and isoleucine and methionine. Mediates the saturable, pH-sensitive and electrogenic cotransport of proline and sodium ions with a stoichiometry of 1:1. May have a role as transporter for neurotransmitter precursors into neurons. In contrast to other members of the neurotransmitter transporter family, does not appear to be chloride-dependent.
Liquid
Biotin
Biotin  Preservative: 0.03% Proclin 300  Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Preservative: 0.03% Proclin 300
Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified  IgG
Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified  IgG  Polyclonal  Sodium-dependent neutral amino acid transporter B(0)AT2 (Sodium- and chloride-dependent neurotransmitter transporter NTT73) (Sodium-coupled branched-chain amino-acid transporter 1) (Solute carrier family 6 member 15)
Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified  IgG  Polyclonal  Sodium-dependent neutral amino acid transporter B(0)AT2 (Sodium- and chloride-dependent neurotransmitter transporter NTT73) (Sodium-coupled branched-chain amino-acid transporter 1) (Solute carrier family 6 member 15) (Transporter v7-3), SLC6A15, B0AT2 NTT73 SBAT1