

🕜 Tel: +1-301-363-4651 🗵 Email: cusabio@cusabio.com 🥃 Website: www.cusabio.com 🧉

STARD10 Antibody

Product Code	CSB-PA896706LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9Y365
Immunogen	Recombinant Human START domain-containing protein 10 protein (1-291AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB; Recommended dilution: WB:1:500-1:2000
Relevance	May play metabolic roles in sperm maturation or fertilization (By similarity). Phospholipid transfer protein that preferentially selects lipid species containing a palmitoyl or stearoyl chain on the sn-1 and an unsaturated fatty acyl chain (18:1 or 18:2) on the sn-2 position. Able to transfer phosphatidylcholine (PC) and phosphatidyetanolamline (PE) between membranes.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	>95%, Protein G purified
lsotype	lgG
Clonality	Polyclonal
Alias	START domain-containing protein 10 (StARD10) (Antigen NY-CO-28) (PCTP- like protein) (PCTP-L) (Serologically defined colon cancer antigen 28) (StAR- related lipid transfer protein 10), STARD10, SDCCAG28
Species	Human
Research Area	Signal Transduction
Target Names	STARD10
Image	Western blot

116kDa	Western blot All lanes: STARD10 antibody at 2µg/ml Lane 1: 293T whole cell lysate
50kDa	Lane 2: k562 whole cell lysate Lane 3: Mouse heart tissue Secondary
35kDa	Goat polyclonal to rabbit IgG at 1/10000 dilution Predicted band size: 33 kDa
25kDa	Observed band size: 33 kDa
20kDa	

1