






## Recombinant Human Large neutral amino acids transporter small subunit 1 (SLC7A5), partial

Storage The street of man and	SB-EP021717HUa3  the shelf life is related to many factors, storage state, buffer ingredients, torage temperature and the stability of the protein itself. Generally, the shelf life if liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.  Form Light Storage Buffer  Storage Buffer  If 59 lyc  Product Type Rollmmunogen Species  Purity G Sequence All Lead Time  13	torage temperature and the stability of the protein itself. Generally, the shelf life f liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 nonths at -20°C/-80°C.
Form Lie Storage Buffer If 59 lye Product Type Re Immunogen Species He Purity G Sequence Al Lead Time 13	01650
Storage Buffer  If 59 lyc Product Type Ref Immunogen Species Purity G Sequence All Lead Time  13	
Product Type Rolling R	iquid or Lyophilized powder
Immunogen Species Ho Purity G Sequence Al Lead Time 13	the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, %-50% glycerol. If the delivery form is lyophilized powder, the buffer before ophilization is Tris/PBS-based buffer, 6% Trehalose.
Purity G Sequence Al Lead Time 13	ecombinant Protein
Sequence Al Lead Time 13	omo sapiens (Human)
Lead Time 13	reater than 90% as determined by SDS-PAGE.
	EEKEEAREKMLAAKSADGSAPAGEGEGVTLQRNI
Research Area Tr	3-23 business days
	ransport
Source E.	.coli
Gene Names SI	LC7A5
Expression Region 16	6-50aa
	epeated freezing and thawing is not recommended. Store working aliquots at °C for up to one week.
Tag Info N	-terminal 6xHis-SUMO-tagged and C-terminal Myc-tagged
Mol. Weight 18	8.0 kDa
Protein Description Pa	artial
Image	(Tris-Glycine gel) Discontinuous SDS-PAGE



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.



## **CUSABIO TECHNOLOGY LLC**





## Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.