



| | |
|--|----------------------------------|
|  | <h2>IgA Polyclonal Antibody</h2> |
|--|----------------------------------|

| | |
|---------------------------------|---|
| Swiss-Prot No.: | P01876/P01877 |
| Molecular Weight: | 37.655 |
| Applications: | WB,IHC,ELISA |
| Reactivity: | Human |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from the Internal region of human IgA. |
| Specificity/Sensitivity: | IgA Polyclonal Antibody detects endogenous levels of IgA protein. |
| Other Names: | IGHA1; Ig alpha-1 chain C region; IGH2; Ig alpha-2 chain C region |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Protein Concentration: | 1 mg/ml |

For Research Use Only

| | |
|-------------------|--|
| |  |
| | Western Blot analysis of various cells using IgA Polyclonal Antibody |
| Gene Name: | IGHA1/IGHA2 |
| Dilution: | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. |