

# Bulk Human IgG2 (IB2) Isotype Control

**Bulk Human IgG2 Isotype Control (IB2)** 

<b>T</b>		<b>n</b>	P 4
Proc	11167	Ken	efits:

11 Todact Benefits.
ichorbio's bulk Human IgG2 Isotype Control (IB2) is manufactured in a cGMP compliant, ISO Quality Standard 9001:2015 facility. ichorbio's low endotoxin antibodies have half the endotoxin of comparable antibodies from <a href="Moleon Bio"><u>Bio Cell</u></a> at less than 1.0 EU/mg. ichorbio: the best antibodies for <i>in vivo</i> research.
Target:
IgG2
Clone:
IB2
Size:
ichorbio's recombinant human IgG2 <i>in vivo</i> antibody is available in the following bulk sizes: 5mg, 25mg, 50mg and 100mg. ichorbio regularly manufactures multi-gram amounts of our human IgG2 antibody - please contact us for pricing.
Isotype:
Human IgG2
Host:
CHO cells
Species Reactivity:
Human
Specificity:
This Human IgG2 isotype control recombinant antibody has been tested against selected species' cells and tissues to assure minimal cross reactivity.
Purification Method:
This monoclonal antibody was purified using multi-step affinity chromatography methods such as Protein A or G depending on the species and isotype.
Background:

ichor.bio // hello@ichor.bio

Recombinant IgG2 human antibody.

Products are for research use only. Not for use in diagnostic or therapeutic procedures. ichorbio, ichorbio logo and all other trademarks are the property of ichorbio LTD © ichorbio 2023 Page 1



# **Concentration:**

>5.0 mg/ml

#### **Formulation:**

This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (PBS) pH 7.2 - 7.4, 150 mM NaCl with no carrier protein, potassium or preservatives added. BSA and azide free

# **Purity:**

>95% by SDS-PAGE and HPLC

>98% by SDS-PAGE and HPLC

#### **Endotoxin:**

? 1.0 EU/mg as determined by the LAL method

? 0.75 EU/mg as determined by the LAL method

#### **Aggregation:**

Aggregation level ? 5%

Aggregation level ? 1%

#### **Storage:**

This antibody is stable for at least 4 weeks when stored at 2-8°C. For long term storage, aliquot in working volumes without diluting and store at  $-20^{\circ}$ C or  $-80^{\circ}$ C. Avoid repeated freeze thaw cycles.

#### **Applications:**

Flow Cytometry, Functional Assays

### **Application Notes:**

Each investigator should determine their own optimal working dilution for specific applications.

## Use:

Products are for research use only.