SARS-CoV-2 Spike RBD (Omicron BA.2.38) Protein





Description	
Source	Recombinant SARS-CoV-2 Spike RBD (Omicron BA.2.38) Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Arg319-Phe541(G339D, S371F, S373P, S375F, T376A, D405N, R408S, K417T, N440K, S477N, T478K, E484A, Q493R, Q498R, N501Y, Y505H).
Accession	QHD43416.1
Molecular Weight	The protein has a predicted MW of 26.07 kDa. Due to glycosylation, the protein migrates to 35-45 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage

Formulation Supplied as 0.22µm filtered solution in PBS (pH 7.4).


Storage Valid for 12 months from date of receipt when stored at -80°C.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The ongoing coronavirus disease 2019 (COVID-19) pandemic has prioritized the development of small-animal models for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The resulting mouse-adapted strain at passage 6 (called MASCp6) showed increased infectivity in mouse lung and led to interstitial pneumonia and inflammatory responses in both young and aged mice after intranasal inoculation. Deep sequencing revealed a panel of adaptive mutations potentially associated with the increased virulence. In particular, the N501Y mutation is located at the receptor binding domain (RBD) of the spike protein.


Assay Data

Tris-Bis PAGE



SARS-CoV-2 Spike RBD (Omicron BA.2.38) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of SARS-CoV-2 Spike RBD (Omicron BA.2.38) is greater than 95% as determined by SEC-HPLC.

Cat. No. BA2-HM138




Assay Data

ELISA Data

SARS-COV-2 Spike RBD (Omicron BA.2.38), His Tag ELISA

0.05μg SARS-COV-2 Spike RBD (Omicron BA.2.38), His Tag Per Well



Immobilized SARS-COV-2 Spike RBD (Omicron BA.2.38) , His Tag at $0.5\mu g/ml$ ($100\mu l/Well$) on the plate. Dose response curve for Human ACE2, hFc Tag with the EC50 of 4.9ng/ml determined by ELISA.