Rhesus macaque CDH17/Cadherin 17 Protein





Description	
Source	Recombinant Rhesus macaque CDH17/Cadherin 17 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Gln23-Thr784.
Accession	A0A1D5R2B4
Molecular Weight	The protein has a predicted MW of 85.60 kDa. Due to glycosylation, the protein migrates to 90-110 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC
Formulation and	l Storage

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in 20mM Tris, 150mM NaCl (pH 8.0). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Liver-intestine cadherin (CDH17) has been known to function as a tumor stimulator and diagnostic marker for almost two decades. In vivo studies showed CDH17 knockout resulted in apoptotic PC tumor death through activating caspase-3 activity. Taken together, CDH17 functions as an oncogenic molecule critical to PC growth by regulating tumor apoptosis signaling pathways and CDH17 could be targeted to develop an anti-PC therapeutic approach.

Assay Data

Tris-Bis PAGE




Rhesus macaque CDH17 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

KAGTUS


Assay Data



The purity of Rhesus macaque CDH17 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Rhesus macaque CDH17, His Tag ELISA 0.05µg Rhesus macaque CDH17, His Tag Per Well



Immobilized Rhesus macaque CDH17, His Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Anti-CDH17 Antibody, hFc Tag with the EC50 of 9.5ng/ml determined by ELISA.