

Human HGFA Protein (pro form)

Cat. No. HGF-HM10A

Description

Source	Recombinant Human HGFA Protein (pro form) is expressed from HEK293 with His tag at the C-Terminus. It contains Gln36-Ser655.
Accession	Q04756
Molecular Weight	The protein has a predicted MW of 67.2 kDa. Due to glycosylation, the protein (pro form) migrates to 85-100 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Hepatocyte growth factor activator (HGFA) is a serine protease initially identified as a potent activator of hepatocyte growth factor/scatter factor. Hepatocyte growth factor/scatter factor is known to be critically involved in tissue morphogenesis, regeneration, and tumor progression, via its receptor, MET. In vivo, HGFA also activates macrophage-stimulating protein, which has roles in macrophage recruitment and inflammatory processes, cellular survival and wound healing through its receptor, RON.

Assay Data


Tris-Bis PAGE



Human HGFA (pro form) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Human HGFA (pro form) is greater than 95% as determined by SEC-HPLC.