

Human CD16a / FCGR3A Protein (176 Val, His & AVI Tag), Biotinylated

Catalog Number: 10389-H27H1-B



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

CD16; CD16A; Fc gamma RIIIa; FCG3; FCGR3; FCGR3III; FCR-10; FCR3III; FCR3IIIA; IGFR3; IMD20

Protein Construction:

A DNA sequence encoding the extracellular domain (Met 1-Gln 208) of human CD16a (AAH17865.1) was fused with a c-terminal polyhistidine tagged AVI tag at the C-terminus. The expressed protein was biotinylated in vivo by the Biotin-Protein ligase (BirA enzyme) which is co-expressed. It is identical to FCGR3A158F/V in the reference.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Biotin/Protein Ratio:

0.7-1 as determined by the HABA assay.

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

1. Measured by its ability to bind recombinant human IgG1 in a functional ELISA.
2. Measured by its binding ability in a functional ELISA. Immobilized human CD16a-AVI-His (Cat:10389-H27H1-B) at 10 µg/ml (100µl/well) can bind recombinant human IgG1 (Fc) (Cat:10702-HNAC) with a linear range of 0.31-5 µg/ml.
3. Using the Octet RED System, the affinity constant (Kd) of CD16a Protein, Human, Recombinant (176 Val, His & AVI Tag), Biotinylated (Cat. 10389-H27H1-B) bound to Human IgG1 was 10⁻⁷-10⁻⁹M.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Gly 17

Molecular Mass:

The secreted recombinant human CD16a consists of 227 amino acids and predicts a molecular mass of 25.6 KDa. By SDS-PAGE under reducing conditions, the apparent molecular mass of the protein is approximately 48 KDa due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

The Fc receptor with low affinity for IgG (FCGR3, or CD16) is encoded by 2 nearly identical genes, FCGR3A and FCGR3B, resulting in tissue-specific expression of alternative membrane-anchored isoforms. FCGR3A, it is also known as CD16a, encodes a transmembrane protein expressed on activated monocytes/macrophages, natural killer (NK) cells, and a subset of T cells. CD16a / FCGR3A is a receptor expressed on NK cells that facilitates antibody dependent cellular cytotoxicity (ADCC) by binding to the Fc portion of various antibodies. CD16a / FCGR3A also has a broader function. CD16a / FCGR3A is directly involved in the lysis of some virus-infected cells and tumor cells by NK cells, independent of antibody binding. Cross-linking of CD16a / FCGR3A on NK cells resulted in increased intracellular Ca²⁺ levels and a cascade of biochemical events similar to those activated by the T cell receptor. CD16a / FCGR3A on human NK cells is a lysis receptor that mediates the direct killing of some virus infected and tumor cells, independent of antibody ligation.

References

3.David Dornan, et al. Effect of FCGR2A and FCGR3A variants on CLL outcome. Blood. 2010 116: 4212-4222

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 ● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 ● Tel:+86-400-890-9989 ● <http://www.sinobiological.com>