

Human EPO Receptor / EPOR Protein (His Tag)

Catalog Number: 10707-H08H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

EPO-R

Protein Construction:

A DNA sequence encoding the extracellular domain (Met 1-Pro 250) of human erythropoietin receptor (NP_000112.1) precursor was fused with a polyhistidine tag at the C-terminus.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Purity: > 98 % as determined by SDS-PAGE

Bio Activity:

Measured by its ability to inhibit EPO-dependent proliferation of TF-1 human erythroleukemic cells. The ED₅₀ for this effect is typically 15-60 ng/mL in the presence of 0.1 U/mL Recombinant Human EPO.

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Ala 25

Molecular Mass:

The secreted recombinant human EPOR consists of 237 amino acids and has a predicted molecular mass of 26.3 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rhEPOR is approximately 34 kDa due to glycosylation.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Erythropoietin (EPO) is the major glycoprotein hormone regulator of mammalian erythropoiesis, and is produced by kidney and liver in an oxygen-dependent manner. The biological effects of EPO are mediated by the specific erythropoietin receptor (EPOR/EPO Receptor) on bone marrow erythroblasts, which transmits signals important for both proliferation and differentiation along the erythroid lineage. EPOR protein is a type α... single-transmembrane cytokine receptor, and belongs to the homodimerizing subclass which functions as ligand-induced or ligand-stabilized homodimers. EPOR signaling prevents neuronal death and ischemic injury. Recent studies have shown that EPO and EPOR protein may be involved in carcinogenesis, angiogenesis, and invasion.

References

- 1.Divoky V, *et al.* (2002) Mouse surviving solely on human erythropoietin receptor (EpoR): model of human EpoR-linked disease. *Blood* 99(10): 3873-4.
- 2.Carruthers SG. (2009) A truncated erythropoietin receptor EPOR-T is associated with hypertension susceptibility. *Clin Pharmacol Ther.* 86(2): 134-6.
- 3.Baltaziak M, *et al.* (2009) Relationships of P53 and Bak with EPO and EPOR in human colorectal cancer. *Anticancer Res.* 29(10):4151-6.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 ● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 ● Tel:+86-400-890-9989 ● <http://www.sinobiological.com>