

# EIF3S1/EIF3J Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological  
Biological Solution Specialist

Catalog Number: 202474-T46

## GENERAL INFORMATION

<b>Immunogen:</b>	E. coli-derived Human EIF3S1/EIF3J fragment
<b>Preparation</b>	Produced in rabbits immunized with E. coli-derived Human EIF3S1/EIF3J fragment, and purified by antigen affinity chromatography.
<b>Ig Type:</b>	Rabbit IgG
<b>Specificity:</b>	Human EIF3S1/EIF3J
<b>Formulation:</b>	PBS, pH7.0 with 0.03% Proclin300
<b>Storage:</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
<b>Alternative Names:</b>	eIF3-alpha,eIF3-p35,EIF3S1

## APPLICATIONS

<b>Applications:</b>	WB,IHC-P,ICC/IF,IP
----------------------	--------------------

## RECOMMENDED CONCENTRATION

<b>IHC-P</b>	IHC-P: 1:100-1:400
<b>ICC/IF</b>	ICC/IF: 1:100-1:500
<b>Western Blot</b>	WB: 1:500-1:2000
<b>Immunoprecipitation</b>	IP:0.5-2µL/mg of lysate


**Please Note: Optimal concentrations/dilutions should be determined by the end user.**

# EIF3S1/EIF3J Antibody, Rabbit PAb, Antigen Affinity Purified




Sino Biological  
Biological Solution Specialist


Catalog Number: 202474-T46



Immunohistochemical staining of human EIF3J in human pancreas with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



Immunofluorescence staining of EIF3J in A431 cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human EIF3J polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Cytoplasm.



Anti-EIF3J rabbit polyclonal antibody at 1:500 dilution

Lane A: HeLa Whole Cell Lysate

Lane B: K562 Whole Cell Lysate

Lane C: Jurkat Whole Cell Lysate


Lane D: U-251MG Whole Cell Lysate

Lysates/proteins at 30 µg per lane.

Secondary  
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique. Performed under reducing conditions.

Predicted band size: 29 kDa  
Observed band size: 34 kDa  
(We are unsure as to the identity of these extra bands.)



EIF3J was immunoprecipitated using:  
Lane A: 0.5 mg HeLa Whole Cell Lysate

4 µL anti-EIF3J rabbit polyclonal antibody and  
60 µg of Immunomagnetic beads Protein  
A/G. Primary antibody:

Anti-EIF3J rabbit polyclonal antibody, at 1:100 dilution

Secondary antibody:  
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution

Developed using the ECL technique.  
Performed under reducing conditions.

Predicted band size: 29 kDa  
Observed band size: 35 kDa