

Mouse PRKAR1A / PRKAR1 / PKR1 Protein (His Tag)

Catalog Number: 51743-M08B



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

1300018C22Rik; R1alpha; Tse-1; Tse1

Protein Construction:

A DNA sequence encoding the mouse PRKAR1A (NP_068680.1) (Met1-Val381) was expressed with a polyhistidine tag at the C-terminus.

Source: Mouse

Expression Host: Baculovirus-Insect Cells

QC Testing

Purity: > 90 % as determined by SDS-PAGE.

Endotoxin:

< 1.0 EU per µg protein as determined by the LAL method.

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Met 1

Molecular Mass:

The recombinant mouse PRKAR1A consists of 392 amino acids and predicts a molecular mass of 44.6 kDa.

Formulation:

Lyophilized from sterile 20 mM Tris, 300 mM NaCl, pH 7.5, 10 % glycerol.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

PRKAR1A, also known as PRKAR1 and PKR1, is one of the regulatory subunits of cAMP-dependent protein kinase A (PKA). PKA can be activated by cAMP. cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its effects by activating PKA, which transduces the signal through phosphorylation of different target proteins. The inactive holoenzyme of PKA is a tetramer composed of two regulatory and two catalytic subunits. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits of PKA have been identified in humans. PRKAR1A was found to be a tissue-specific extinguisher that down-regulates the expression of seven liver genes in hepatoma x fibroblast hybrids. Three alternatively spliced transcript variants encoding the same protein have been observed.

References

1. Huang L J. et al., 1997, Proc Natl Acad Sci. 94 (21): 11184-9. 2. Herberg F W. et al., 2000, J Mol Biol. 298 (2): 329-39. 3. Scambler P. et al., 1987, Am J Hum Genet. 41 (5): 925-32.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 • Tel:+86-400-890-9989 • <http://www.sinobiological.com>