Rat KIM-1 / TIM1 / HACVR1 Protein (His Tag)

Catalog Number: 80003-R07H



General Information

Gene Name Synonym:

HAVCR1

Protein Construction:

A DNA sequence encoding the extracellular domain of rat HAVCR1 (NP_775172.1) (Ser 18-Val 238) was expressed, fused with a polyhistidine tag at the N-terminus.

Source: Rat

Expression Host: HEK293 Cells

QC Testing

Purity: > 97 % as determined by SDS-PAGE

Endotoxin:

< 1.0 EU per µg of the protein as determined by the LAL method

Stability:

Samples are stable for up to twelve months from date of receipt $\,$ at -70 $\,$ $^{\circ}$ C

Predicted N terminal: His

Molecular Mass:

The recombinant rat HAVCR1 comprises 237 amino acids and predicts a molecular mass of 26.7 kDa. As a result of glycosylation, the apparent molecular mass of the rat HAVCR1 is approximately 55-65 kDa in SDS-PAGE under reducing conditions.

Formulation:

Lyophilized from sterile PBS, pH 7.4

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide


Storage:

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

HAV cellular receptor 1 (HAVCR1), also known as Kidney injury molecule 1 (KIM-1) and T cell immunoglobulinmucin 1 (TIM-1), is a type â... integral membrane glycoprotein. KIM-1 protein is widely expressed with highest levels in kidney and testis. It has been shown to play a major role as a human susceptibility gene for asthma, allergy and autoimmunity. IgA1lambda is a specific ligand of KIM-1 protein and that their association has a synergistic effect in virus-receptor interactions. KIM-1 involves in the pathogenesis of acute kidney injury. It had been confirmed that KIM-1 is a human urinary renal dysfunction biomarker. Moreover, KIM-1 protein is a novel regulatory molecule of flow-induced calcium signaling.

References

1.Tami C, et al. (2007) Immunoglobulin A (IgA) is a natural ligand of hepatitis A virus cellular receptor 1 (HAVCR1), and the association of IgA with HAVCR1 enhances virus-receptor interactions. J Virol. 81(7): 3437-46. 2.Rees AJ, et al. (2008) Kim-1/Tim-1: from biomarker to therapeutic target? Nephrol Dial Transplant. 23(11): 3394-6. 3.Chaturvedi S, et al. (2009) Assay validation for KIM-1: human urinary renal dysfunction biomarker. Int J Biol Sci. 5(2): 128-34.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 • Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288
■ Tel:+86-400-890-9989
■ http://www.sinobiological.com