Human NCR2 Gene cDNA clone plasmid

Catalog Number: HG11550-M



General Information

Gene: natural cytotoxicity triggering receptor 2.

Official Symbol: NCR2

Synonym: LY95, CD336, NKP44, NK-p44,

dJ149M18.1, NCR2

Source: Human

cDNA Size: 831bp

RefSeq: NM_004828.3

Plasmid: pMD-NCR2

Description

Lot: Please refer to the label on the tube

Sequence Description:

Identical with the Gene Bank Ref. ID sequence.

Vector:

pMD18-T

Shipping carrier:

Each tube contains approximately 10 µg of lyophilized plasmid.

Storage:

The lyophilized plasmid can be stored at ambient temperature for three months.

Quality control:

The plasmid is confirmed by full-length sequencing with primers in the sequencing primer list.

Sequencing primer list:

5' GCCAGGGTTTTCCCAGTCACGAC 3' M13-47:

RV-M: 5' GAGCGGATAACAATTTCACACAGG 3'

Other M13 primers can also be used as sequencing primers.

Plasmid Resuspension protocol

- 1. Centrifuge at 5,000×g for 5 min.
- 2. Carefully open the tube and add 100 µl of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than
- 5. Store the plasmid at -20 °C.

The plasmid is ready for:

- Restriction enzyme digestion
- PCR amplification
- · E. coli transformation
- DNA sequencing

E.coli strains for transformation (recommended but not limited)

Most commercially available competent cells are appropriate for the plasmid, e.g. TOP10, DH5α and TOP10F'.

Human NCR2 Gene cDNA clone plasmid

Catalog Number: HG11550-M



Vector Information

pMD18-T Vector is a high-efficiency TA cloning vector constructed from pUC18, of which multiple cloning sites as shown below. The pMD18-T Vector is 2.6kb in size and contains the amplicin resistance gene for selection. The coding sequence was inserted by TA cloning at site 425.

Physical Map of pMD18 (MCS destroyed):

