# Human B3GALT4 ORF mammalian expression plasmid, N-Myc tag



#### Catalog Number: HG14748-NM

#### **General Information**

Gene :UDP-Gal:betaGlcNAcbeta1,3-galactosyltransferase, polypeptide 41Official Symbol :B3GALT4Synonym :GALT2, GALT4, BETA3GALT4Source :HumancDNA Size:1137bpRefSeq :BC032574

#### Description

Lot : Please refer to the label on the tube

Vector : pCMV3-SP-N-Myc

#### Shipping carrier :

Each tube contains approximately 10 µg of lyophilized plasmid.

#### Storage :

The lyophilized plasmid can be stored at ambient temperature for three months.

#### **Quality control :**

The plasmid is confirmed by full-length sequencing with primers in the sequencing primer list.

#### Sequencing primer list :

| pCMV3-F:     | 5' CAGGTGTCCACTCCCAGGTCCAAG 3' |  |
|--------------|--------------------------------|--|
| pcDNA3-R :   | 5' GGCAACTAGAAGGCACAGTCGAGG 3' |  |
| Or           |                                |  |
| Forward T7 : | 5' TAATACGACTCACTATAGGG 3'     |  |
| ReverseBGH : | 5' TAGAAGGCACAGTCGAGG 3'       |  |

pCMV3-F and pcDNA3-R are designed by Sino Biological Inc. Customers can order the primer pair from any oligonucleotide supplier.

#### **Plasmid Resuspension protocol**

- 1. Centrifuge at  $5,000 \times g$  for 5 min.
- 2. Carefully open the tube and add 100  $\mu$ l of sterile water to dissolve the DNA.
- Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than  $5000 \times g$ .
- 5. Store the plasmid at -20  $^{\circ}$ C.

#### The plasmid is ready for:

- Restriction enzyme digestion
- PCR amplification
- E. coli transformation
- DNA sequencing

## *E.coli* strains for transformation (recommended but not limited)

Most commercially available competent cells are appropriate for the plasmid, e.g. TOP10, DH5  $\alpha$  and TOP10F  $\dot{}$  .

## Human B3GALT4 ORF mammalian expression plasmid, N-Myc tag



Catalog Number: HG14748-NM

#### **Vector Information**

All of the pCMV vectors are designed for high-level stable and transient expression in mammalian hosts. High-level stable and non-replicative transient expression can be carried out in most mammalian cells. The vectors contain the following elements:

•Human enhanced cytomegalovirus immediate-early (CMV) promoter for high-level expression in a wide range of mammalian cells.

• Hygromycin resistance gene for selection of mammalian cell lines.

• A Kozak consensus sequence to enhance mammalian expression.


| Vector Name                     | pCMV3-SP-N-Myc                  |
|---------------------------------|---------------------------------|
| Vector Size                     | 6149bp                          |
| Vector Type                     | Mammalian Expression Vector     |
| Expression Method               | Constiutive, Stable / Transient |
| Promoter                        | CMV                             |
| AntibioticResistance            | Kanamycin                       |
| Selection In<br>Mammalian Cells | Hygromycin                      |
| Protein Tag                     | Mvc                             |

### pCMV3-SP-N-Myc (suitable for secretary

and membane protein expession)



#### **Physical Map**




#### Comments for pCMV3-SP-N-Myc:

CMV promoter: bases 250-837 enhancer: bases 838-1445 SV40 early promoter: bases 2390-2759 Hygromycin ORF: bases 2777-3802 pUC origin: bases 4445-5118 Kanamycin ORF: bases 5192-6007

#### Description

| Vector Name                  | pCMV3-SP-N-Myc  |
|------------------------------|---|
| Vector Size                  | 6149bp  |
| Vector Type                  | Mammalian Expression Vector   |
| Expression Method            | Constitutive, Stable / Transient                                    |
| Promoter                     | CMV   |
| Antibiotic Resistance        | Kanamycin   |
| Selection In Mammalian Cells | Hygromycin  |
| Protein Tag                  | Мус   |
| Sequencing Primer            | Forward:T7(TAATACGACTCACTATAGGG)<br>Reverse:BGH(TAGAAGGCACAGTCGAGG) |

#### Schematic of pCMV3-SP-N-Myc Multiple Cloning Sites



pCMV3-SP-N-Myc is recommended for constructing the N-Myc tag secretary and membrane proteins expression vector which containing a naïve signal peptide. An universal signal peptide is used to instead the naïve signal peptide.