Human RIC8A ORF mammalian expression plasmid, N-Flag tag



Catalog Number: HG15892-NF

General Information

Gene: resistance to inhibitors of cholinesterase

8 homolog A (C. elegans)

Official Symbol: RIC8A

RIC8 Synonym:

Source: Human

cDNA Size: 1593bp

RefSeq: BC121808

Description

Lot: Please refer to the label on the tube

Vector: pCMV3-SP-N-FLAG

Shipping carrier:

Each tube contains approximately 10 µg of lyophilized plasmid.

Storage:

The lyophilized plasmid can be stored at ambient temperature for three months.

Quality control:

The plasmid is confirmed by full-length sequencing with primers in the sequencing primer list.

Sequencing primer list:

pCMV3-F: 5' CAGGTGTCCACTCCCAGGTCCAAG 3' 5' GGCAACTAGAAGGCACAGTCGAGG 3'

pcDNA3-R:

Or

Forward T7: 5' TAATACGACTCACTATAGGG 3'

ReverseBGH: 5' TAGAAGGCACAGTCGAGG 3'

pCMV3-F and pcDNA3-R are designed by Sino Biological Inc. Customers can order the primer pair from any oligonucleotide supplier.

Plasmid Resuspension protocol

- Centrifuge at $5,000 \times g$ for 5 min.
- Carefully open the tube and add 100 µl of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000×g.
- 5. Store the plasmid at -20 °C.

The plasmid is ready for:

- Restriction enzyme digestion
- PCR amplification
- · E. coli transformation
- DNA sequencing

E.coli strains for transformation (recommended but not limited)

Most commercially available competent cells are appropriate for the plasmid, e.g. TOP10, DH5α and TOP10F'.

Human RIC8A ORF mammalian expression plasmid, N-Flag tag



Catalog Number: HG15892-NF

Vector Information

All of the pCMV vectors are designed for high-level stable and transient expression in mammalian hosts. High-level stable and non-replicative transient expression can be carried out in most mammalian cells. The vectors contain the following elements:

•Human enhanced cytomegalovirus immediate-early (CMV) promoter for high-level expression in a wide range of mammalian cells.

· Hygromycin resistance gene for selection of mammalian cell lines.

 A Kozak consensus sequence to enhance mammalian expression.

Vector Name pCMV3-SP-N-FLAG

Vector Size 6143bp

Vector Type Mammalian Expression Vector Expression Method Constitutive, Stable / Transient

Hygromycin

Promoter

Antibiotic Resistance Kanamycin

Selection In Mammalian Cells


Protein Tag

FLAG

pCMV3-SP-N-FLAG (suitable for secretary and membane protein expession)



Physical Map



Comments for pCMV3-SP-N-FLAG:


CMV promoter: bases 250-837 enhancer: bases 838-1445

SV40 early promoter: bases 2384-2753 Hygromycin ORF: bases 2771-3793 pUC origin: bases 4439-5112 Kanamycin ORF: bases 5186-6001

Description

Vector Name	pCMV3-SP-N-FLAG
Vector Size	6143bp
Vector Type	Mammalian Expression Vector
Expression Method	Constitutive, Stable / Transient
Promoter	CMV
Antibiotic Resistance	Kanamycin
Selection In Mammalian Cells	Hygromycin
Protein Tag	FLAG
Sequencing Primer	Forward:T7(TAATACGACTCACTATAGGG) Reverse:BGH(TAGAAGGCACAGTCGAGG)

Schematic of pCMV3-SP-N-FLAG Multiple Cloning Sites



pCMV3-SP-N-Flag is recommended for constructing the N-FLAG tag secretary and membrane proteins expression vector which containing a naïve signal peptide. An universal signal peptide is used to instead the naïve signal peptide.