



Sino Biological
Biological Solution Specialist

Anti-SSR3 Magnetic Beads Immunoprecipitation (IP) Kit

Catalog Number: MB202607-T46

Please read this instruction manual carefully before using the product

Product Contents

| | Contents | Package 1 (20 Tests) | Package 2 (100 Tests) | Storage |
|---|---|---|-----------------------|---------------------|
| 1 | Anti-SSR3 Magnetic Beads ^{1 3} | 1 mL | 5 mL | 2-8°C for 12 months |
| 2 | NP40 Cell Lysis Buffer ² | 4 mL | 22 mL | -20°C for 12 months |
| 3 | 5×TBST (pH7.4) | Required but not supplied | | |
| 4 | 1×TBST (pH7.4) | Required but not supplied | | |
| 5 | ddH ₂ O | Required but not supplied | | |
| 6 | Alkaline Elution Buffer | 3 mL | 15 mL | 2-8°C for 12 months |
| 7 | Acidity Elution Buffer | 3 mL | 15 mL | 2-8°C for 12 months |
| 8 | Neutralization Buffer | 2 mL | 8 mL | 2-8°C for 12 months |
| 9 | Magnetic Separator | One Simple Magnetic Separator (Cat# MAGS001) | | |

[1] The IP KIT contains anti-SSR3 Immunomagnetic Beads(2 mg/mL) in phosphate buffered saline (PBS, pH 7.4) with sodium azide (0.1%).

[2] Using NP-40 cell lysate buffer in the kit is required, otherwise, the magnetic beads may be precipitated.

[3] Immunomagnetic Beads kits are shipped at ambient temperature in which immunomagnetic beads are provided in liquid buffer.

Product Description

The Anti-SSR3 Immunomagnetic Beads, conjugated with Anti-SSR3 antibody, are used for immunoprecipitation (IP) of SSR3 proteins which expressed in vitro expression systems and bacterial and mammalian cell lysates.

For IP, the beads are added to a sample containing SSR3 proteins to form a bead-protein complex. The complex is removed from the solution manually using a Magnetic Separator. The bound SSR3 proteins are dissociated from the Immunomagnetic Beads using an elution buffer.

Antibody Information

Antibody: SSR3 Antibody, Rabbit PAb, Antigen Affinity Purified (Cat# 202607-T46)

Immunogen: E. coli-derived Human SSR3 fragment

Isotype: Rabbit IgG

Specificity: Human SSR3

Preparation: Produced in rabbits immunized with E. coli-derived Human SSR3 fragment, and purified by antigen affinity chromatography.

Applications: IP, Minimum Protein Purification

Alternative Names: SSR3




Fig. 1 Immunoprecipitation (IP) Protocol

Protocol

The protocol (Fig. 1) uses 50 μL Anti-SSR3 Immunomagnetic Beads, but this can be scaled up or down as required.

Cell Lysis

Cells may be lysed using any standard cell lysis protocol in accordance with your starting materials. **We suggest using NP40 Cell Lysis Buffer (supplied with kit).**

Immunoprecipitate Target Antigen

1. Add 50 μL of Immunomagnetic Beads into a 1.5 mL microcentrifuge tube.
2. Add 150 μL of $1\times$ TBST buffer to the Immunomagnetic Beads and gently vortex to mix.
3. Place the tube into a Magnetic Separator to collect the beads against the wall side of the tube. Remove and discard the supernatant.
4. Add 1 mL of $1\times$ TBST buffer to the tube. Invert the tube several times or gently vortex to mix for 1 min. Collect Immunomagnetic Beads with a Magnetic Separator. Remove and discard the supernatant.
5. Add the sample containing target protein (Cell lysate: 0.5-1mg; Recombinant protein: 5-25 μg) to the pre-washed Immunomagnetic Beads, add $1\times$ TBST buffer until final volume to 200-500 μL , and incubate at 37°C for 20-30 min (or at room temperature for 2-3h) with mixing.
6. Collect the Immunomagnetic Beads with a Magnetic Separator, remove the unbound sample and save for analysis.
7. Add 300 μL of $5\times$ TBST buffer to the tube and gently mix. Collect the Immunomagnetic Beads and discard the supernatant. Repeat this wash twice.
8. Add 300 μL of ddH_2O to the tube and gently mix. Collect the Immunomagnetic Beads on a Magnetic Separator and discard the supernatant.

Elute Target Antigen.

A. Alkaline Elution

1. Add 100 μL of Alkaline Elution buffer to the tube.
2. Gently vortex to mix and incubate the sample at room temperature on a rotator for 5 min.
3. Magnetically separate the Immunomagnetic Beads and save the supernatant containing the target antigen.
4. To neutralize the sample, add 50 μL of Neutralization Buffer for each 100 μL of eluate.

B. Acidity Elution

1. Add 100 μL Acidity Elution Buffer.
2. Gently vortex to mix and incubate the sample at room temperature on a rotator for 5-10 min.
3. Magnetically separate the Immunomagnetic Beads and save the supernatant containing the target antigen.
4. To neutralize the low pH, add 15 μL of Neutralization Buffer for each 100 μL of eluate.

C. Denaturing Elution

1. Add 10 μL of $2\times$ SDS-PAGE Sample Loading Buffer to the tube.
2. Gently vortex to mix and incubate the sample at $95-100^\circ\text{C}$ for 5-10 min.
3. Magnetically separate the Immunomagnetic Beads and save the supernatant containing the antigen.

General Test System of Sino Biological Inc. (for reference) :

| | Recombinant Protein | Cell Lysate |
|----------------|--|-------------|
| Sample Quality | 10 μg add into 0.5mg cell lysate (without interfering proteins) | 0.5mg |
| Final Volume | 300 μL | |
| Incubate Time | Room temperature, 2h | |
| Elute | Using 10 μL of $2\times$ SDS-PAGE Sample Loading Buffer | |

Reference Information

Related Products

| Products | Cat No. |
|--|----------|
| Magnetic Separator-1.5 (2 tubes) | MAGS001 |
| Immunoprecipitation Kit -Immunomagnetic Beads Protein A Kit | BA10600 |
| Immunoprecipitation Kit -Immunomagnetic Beads Protein G Kit | BG13103 |
| Immunoprecipitation Kit -Immunomagnetic Beads Protein L Kit | BL11044 |
| Immunoprecipitation Kit -Immunomagnetic Beads Protein A/G Kit | BAG001 |
| Immunoprecipitation Kit -Anti-DYKDDDDK(Flag®) Tag Immunomagnetic Beads Kit | TB101274 |
| Immunoprecipitation Kit -Anti-GFP Tag Immunomagnetic Beads Kit | TB13105 |
| Immunoprecipitation Kit -Anti-Myc Tag Immunomagnetic Beads Kit | TB100029 |
| Immunoprecipitation Kit -Anti-HA Tag Immunomagnetic Beads Kit | TB100028 |
| Immunoprecipitation Kit -Anti-V5 Tag Immunomagnetic Beads Kit | TB100378 |
| Immunoprecipitation Kit -Anti-GST Tag Immunomagnetic Beads Kit | TB11213 |
| Magpoins™ His-Tag Immunoprecipitation Kit | TBN001 |

Trouble Shooting

| Problem | Possible Cause | Solution |
|----------------------------------|-------------------------------------|---|
| Little or no protein is detected | Protein degraded | Include protease inhibitors (PMSF) in the lysis buffer |
| | | Use new lysate or lysate stored at -80° C |
| | No or minimal protein was expressed | Verify protein expression by SDS-PAGE or Western blot |
| | | Analysis of the lysate using an positive control as a reference |

| Problem | Possible Cause | Solution | |
|--------------------------------------|--|--|---|
| Little or no protein is detected | No or minimal protein was expressed | Increase the amount of lysate used for IP/Co-IP | |
| | | Use a more sensitive detection system | |
| Magnetic Beads aggregated | Magnetic Beads were frozen or centrifuged | Handle the Beads as directed in the instructions | |
| | Buffer was incompatible with magnetic beads | | |
| | Detergent was not added to the wash and bind solutions | | |
| Failure to co-IP interacting protein | Wash conditions were too stringent for the weak or transient interaction | Reduce the number of washes | |
| | | Lower the ionic strength of the wash buffer | |
| | Interacting protein was expressed at a low level | Apply additional protein sample | |
| | | Use a more sensitive detection system | |
| | Buffer system was not optimal for the protein: protein interaction | Insufficient sample was loaded on the gel for Western blot detection | Optimize the co-IP buffer |
| | | | Elute sample in 30% acetonitrile 0.5% formic acid, then Bring the sample back up in SDS-PAGE Sample Loading Buffer and load entire elution fraction on |