



Rat Anti-Human IL-4

Cat. No.	Format	Size
10117-01	Purified (UNLB)	0.5 mg
10117-08	Biotin (BIOT)	0.5 mg
10117-14	Low Endotoxin, Azide-Free (LE/AF)	0.5 mg

Overview

Clone	MP4-25D2
Isotype	Rat IgG _{1κ}
Immunogen	CHO-expressed human IL-4
Specificity	Human/Rhesus/Cynomolgus/Baboon IL-4
Alternate Name(s)	Interleukin-4, T cell growth factor-2, TCGF-2, Ia inducing factor, IaIF, B cell stimulating factor-1, BSF-1, Hodgkin's cell growth factor, HCGF, mast cell growth factor-2, MCGF-2, macrophage fusion factor, MFF, T cell growth factor-2, TCGF-2, Hodgkin's cell growth factor, HCGF

Applications

ELISA-Detection – Quality tested ^{1,2,4,5}
ELISA-Capture – Reported in literature ³
ELISPOT-Capture – Reported in literature ^{2,6}
ELISPOT-Detection – Reported in literature ⁷
FC – Reported in literature ⁸⁻¹⁰
IHC-FS – Reported in literature ¹¹⁻¹³
Neut – Reported in literature ^{1,14,15}
Multiplex-Detection – Reported in literature ¹⁶

Working Dilutions

ELISA	BIOT conjugate	1:2,000 – 1:4,000
--------------	----------------	-------------------

Other Applications Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The low endotoxin, azide-free (LE/AF) antibody is supplied as 0.5 mg purified immunoglobulin in 1.0 mL of PBS. Contains no preservative; handle under aseptic conditions. Store at 2-8°C or aliquot into smaller volumes and store at -20°C. Avoid multiple freeze / thaw cycles.
- Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific SDS.

References

1. Abrams JS, Roncarolo M, Yssel H, Andersson U, Gleich GJ, Silver JE. Strategies of anti-cytokine monoclonal antibody development: immunoassay of IL-10 and IL-5 in clinical samples. *Immunol Rev.* 1992;127:5-24. (ELISA-Detection, Neut)
2. Mahanty S, Abrams JS, King CL, Limaye AP, Nutman TB. Parallel regulation of IL-4 and IL-5 in human helminth infections. *J Immunol.* 1992;148:3567-71. (ELISA-Detection, ELISPOT-Capture)
3. Nyindo M, Kariuki TM, Mola PW, Farah IO, Elson L, Blanton RE, et al. Role of adult worm antigen-specific immunoglobulin E in acquired immunity to *Schistosoma mansoni* infection in baboons. *Infect Immun.* 1999;67:636-42. (ELISA-Capture, Baboon Reactivity)
4. Abrams JS. Immunozytometric assay of mouse and human cytokines using NIP-labeled anti-cytokine antibodies. *Curr Protoc Immunol.* 2001;6.20:1-15. (ELISA-Detection)
5. Zhao W, Oskeritzian CA, Pozez AL, Schwartz LB. Cytokine production by skin-derived mast cells: endogenous proteases are responsible for degradation of cytokines. *J Immunol.* 2005;175:2635-42. (ELISA-Detection)
6. Klinman D. ELISPOT assay to detect cytokine-secreting murine and human cells. *Curr Protoc Immunol.* 2008;6.19:1-9. (ELISPOT-Capture)
7. Guerkov RE, Targoni OS, Kreher CR, Boehm BO, Herrera MT, Tary-Lehmann M, et al. Detection of low-frequency antigen-specific IL-10-producing CD4⁺ T cells via ELISPOT in PBMC: cognate vs. nonspecific production of the cytokine. *J Immunol Methods.* 2003;279:111-21. (ELISPOT-Detection)
8. Koshiba M, Rosin DL, Hayashi N, Linden J, Sitkovsky MV. Patterns of A_{2A} extracellular adenosine receptor expression in different functional subsets of human peripheral T cells. Flow cytometry studies with anti-A_{2A} receptor monoclonal antibodies. *Mol Pharmacol.* 1999;55:614-24. (FC)
9. Bandeira-Melo C, Sugiyama K, Woods LJ, Weller PF. Cutting edge: eotaxin elicits rapid vesicular transport-mediated release of preformed IL-4 from human eosinophils. *J Immunol.* 2001;166:4813-7. (FC)
10. Gansuud B, Goodwin J, Asiedu CK, Jiang XL, Jargal U, Andrades P, et al. Invariant natural killer T cells from rhesus macaque spleen and peripheral blood are phenotypically and functionally distinct populations. *J Med Primatol.* 2008;37:1-11. (FC, Rhesus Reactivity)
11. Andersson J, Abrams J, Björk L, Funa K, Litton M, Ågren K, et al. Concomitant in vivo production of 19 different cytokines in human tonsils. *Immunology.* 1994;83:16-24. (IHC-FS)
12. Ulfgren A, Lindblad S, Klareskog L, Andersson J, Andersson U. Detection of cytokine producing cells in the synovial membrane from patients with rheumatoid arthritis. *Ann Rheum Dis.* 1995;54:654-61. (IHC-FS)
13. Sparrelid E, Emanuel D, Fehniger T, Andersson U, Andersson J. Interstitial pneumonitis in bone marrow transplant recipients is associated with local production of TH2-type cytokines and lack of T cell-mediated cytotoxicity. *Transplantation.* 1997;63:1782-9. (IHC-FS)
14. Punnonen J, Yssel H, de Vries JE. The relative contribution of IL-4 and IL-13 to human IgE synthesis induced by activated CD4⁺ or CD8⁺ T cells. *J Allergy Clin Immunol.* 1997;100:792-801. (Neut)
15. Cousins DJ, Lee TH, Staynov DZ. Cytokine coexpression during human Th1/Th2 cell differentiation: direct evidence for coordinated expression of Th2 cytokines. *J Immunol.* 2002;169:2498-506. (Neut)
16. Hutchinson KL, Villinger F, Miranda ME, Ksiazek TG, Peters CJ, Rollin PE. Multiplex analysis of cytokines in the blood of cynomolgus macaques naturally infected with Ebola virus (Reston serotype). *J Med Virol.* 2001;65:561-6. (Multiplex-Detection, Cynomolgus Reactivity)

TB10117
08-Oct-21

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA **Mailing Address:** P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • **Fax:** 205.945.8768

Email: info@southernbiotech.com • **Website:** www.southernbiotech.com