




Goat Anti-Human Kappa

Cat. No.	Format	Size
2060-01	Purified (UNLB)	1.0 mg
2060-02	Fluorescein (FITC)	1.0 mg
2060-03	Rhodamine (TRITC)	1.0 mg
2060-04	Alkaline Phosphatase (AP)	1.0 mL
2060-05	Horseradish Peroxidase (HRP)	1.0 mL
2060-06	β -galactosidase (BGAL)	1.0 mL
2060-07	Texas Red [®] (TXRD)	1.0 mg
2060-08	Biotin (BIOT)	1.0 mg
2060-09	R-phycoerythrin (PE)	0.5 mg
2060-30	Alexa Fluor [®] 488 (AF488)	1.0 mg
2060-31	Alexa Fluor [®] 647 (AF647)	1.0 mg
2060-32	Alexa Fluor [®] 555 (AF555)	1.0 mg



FLISA plate was coated with purified human IgG κ , IgM κ , IgA κ , IgG λ , IgM λ , and IgA λ . Immunoglobulins were detected with serially diluted Goat Anti-Human Kappa-FITC (SB Cat. No. 2060-02).

Description

Specificity	Reacts with human κ light chains
Source	Pooled antisera from goats hyperimmunized with human κ light chains
Cross Adsorption	Human λ light chains; may react with κ light chains from other species
Purification	Affinity chromatography on human κ light chains covalently linked to agarose

Applications

Quality tested applications include –

ELISA¹⁻⁹
 FLISA
 FC¹⁹⁻²²

Other referenced applications include –

ELISPOT¹⁰
 IHC-PS¹¹
 ICC¹²
 WB^{4,13-17}
 IP¹⁸
 SPR²³
 Purification²⁴
 Stim²⁵

Working Dilutions

ELISA	AP conjugate	1:2,000 – 1:4,000
	HRP conjugate	1:4,000 – 1:8,000
	BGAL conjugate	1:500
	BIOT conjugate	1:5,000 – 1:20,000
FLISA	FITC, TRITC, TXRD, AF488, and AF555 conjugates	1:100 – 1:400
	PE and AF647 conjugates	$\leq 1 \mu\text{g/mL}$
Flow Cytometry	FITC, BIOT, and AF488 conjugates	$\leq 1 \mu\text{g}/10^6 \text{ cells}$
	PE and AF647 conjugates	$\leq 0.1 \mu\text{g}/10^6 \text{ cells}$
	For flow cytometry, the suggested use of these reagents is in a final volume of 100 μL	
Other Applications	Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.	

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- The fluorescein (FITC), rhodamine (TRITC), Texas Red® (TXRD), Alexa Fluor® 488 (AF488), Alexa Fluor® 555 (AF555), and Alexa Fluor® 647 (AF647) conjugates are supplied as 1.0 mg in 1.0 mL of PBS/NaN₃. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL of stock solution in 50 mM Tris/1 mM MgCl₂/50% glycerol, pH 8.0, containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The β-galactosidase (BGAL) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50% PBS containing NaN₃ as preservative. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL of PBS/NaN₃. Store at 2-8°C.
- The R-phycoerythrin (PE) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

Warning

Some reagents contain sodium azide. Please refer to product specific (M)SDS.

References

1. Pancook JD, Beuerlein G, Pecht G, Tang Y, Nie Y, Wu H, et al. In vitro affinity maturation of human IgM antibodies reactive with tumor-associated antigens. *Hybrid Hybridomics*. 2001;20:383-96. (ELISA)
2. Akamatsu Y, Pakabunto K, Xu Z, Zhang Y, Tsurushita N. Whole IgG surface display on mammalian cells: Application to isolation of neutralizing chicken monoclonal anti-IL-12 antibodies. *J Immunol Methods*. 2007;327:40-52. (ELISA)
3. Guan Y, Sajadi MM, Kamin-Lewis R, Fouts TR, Dimitrov A, Zhang Z, et al. Discordant memory B cell and circulating anti-Env antibody responses in HIV-1 infection. *Proc Natl Acad Sci USA*. 2009;106:3952-7. (ELISA)
4. Lai H, Engle M, Fuchs A, Keller T, Johnson S, Gorlatov S, et al. Monoclonal antibody produced in plants efficiently treats West Nile virus infection in mice. *Proc Natl Acad Sci USA*. 2010;107:2419-24. (ELISA, WB)
5. Boross P, Lohse S, Nederend M, Jansen JH, van Tetering G, Dechant M, et al. IgA EGFR antibodies mediate tumour killing in vivo. *EMBO Mol Med*. 2013;5:1213-26. (ELISA)
6. Kehoe JW, Whitaker B, Bethea D, Lacy ER, Boakye K, Santulli-Marotto S, et al. Isolation and optimization for affinity and biophysical characteristics of anti-CCL17 antibodies from the V_H1-69 germ-line gene. *Protein Eng Des Sel*. 2014;27:199-206. (ELISA)
7. Wyrzucki A, Dreyfus C, Kohler I, Steck M, Wilson IA, Hangartner L. Alternative recognition of the conserved stem epitope in influenza A virus hemagglutinin by a V_H3-30-encoded heterosubtypic antibody. *J Virol*. 2014;88:7083-92. (ELISA)
8. Nagano R, Masuda K. Establishment of a signal peptide with cross-species compatibility for functional antibody expression in both Escherichia coli and Chinese hamster ovary cells. *Biochem Biophys Res Commun*. 2014;447:655-9. (ELISA)
9. London AS, Mackay K, Lihon M, He Y. Gel filtration chromatography as a method for removing bacterial endotoxin from antibody preparations. *Biotechnol Prog*. 2014;30:1497-1501. (ELISA)
10. Saletti G, Çuburu N, Yang JS, Dey A, Czerkinsky C. Enzyme-linked immunospot assays for direct ex vivo measurement of vaccine-induced human humoral immune responses in blood. *Nat Protoc*. 2013;8:1073-87. (ELISPOT)
11. Bonda DJ, Webber KM, Siedlak SL, Perry G, Friedland RP, Smith MA. The pathology of Alzheimer disease elicits an in vivo immunological response. *Am J Immunol*. 2007;3:10-14. (IHC-PS)
12. Campana D, Farrant J, Inamdar N, Webster AD, Janosy G. Phenotypic features and proliferative activity of B cell progenitors in X-linked agammaglobulinemia. *J Immunol*. 1990;145:1675-80. (ICC)
13. Singh G, Parker S, Hobart P. The development of a bicistronic plasmid DNA vaccine for B-cell lymphoma. *Vaccine*. 2002;20:1400-11. (WB)
14. Corisdeo S, Wang B. Functional expression and display of an antibody Fab fragment in Escherichia coli: study of vector designs and culture conditions. *Protein Expr Purif*. 2004;34:270-9. (WB)
15. Phoolcharoen W, Bhoo SH, Lai H, Ma J, Arntzen CJ, Chen Q, et al. Expression of an immunogenic Ebola immune complex in Nicotiana benthamiana. *Plant Biotechnol J*. 2011;9:807-16. (WB)
16. He J, Peng L, Lai H, Hurtado J, Stahnke J, Chen Q. A plant-produced antigen elicits potent immune responses against West Nile virus in mice. *Biomed Res Int*. 2014;2014:952865. (WB)
17. Helman D, Toister-Achituv M, Bar-Shimon M, Cohen B, Otmir I, Berger N, et al. Novel membrane-bound reporter molecule for sorting high producer cells by flow cytometry. *Cytometry A*. 2014;85:162-8. (WB)
18. Santiago T, Kulemzin SV, Reshetnikova ES, Chikaev NA, Volkova OY, Mechetina LV, et al. FCRLA is a resident endoplasmic reticulum protein that associates with intracellular Igs, IgM, IgG and IgA. *Int Immunol*. 2011;23:43-53. (IP)
19. Minegishi Y, Conley ME. Negative selection at the pre-BCR checkpoint elicited by human μ heavy chains with unusual CDR3 regions. *Immunity*. 2001;14:631-41. (FC)
20. Hinton PR, Xiong JM, Johlfis MG, Tang MT, Keller S, Tsurushita N. An engineered human IgG1 antibody with longer serum half-life. *J Immunol*. 2006;176:346-56. (FC)
21. Dong J, Sereno A, Aivazian D, Langley E, Miller BR, Snyder WB, et al. A stable IgG-like bispecific antibody targeting the epidermal growth factor receptor and the type I insulin-like growth factor receptor demonstrates superior anti-tumor activity. *mAbs*. 2011;3:273-88. (FC)
22. Xiang Y, Park S, Garrard WT. A major deletion in the V_κ-J_κ intervening region results in hyper-elevated transcription of proximal V_κ genes and a severely restricted repertoire. *J Immunol*. 2014;193:3746-54. (FC)
23. Wagner K, Kwakkenbos MJ, Claassen YB, Majoor K, Böhne M, van der Sluijs KF, et al. Bispecific antibody generated with sortase and click chemistry has broad anti-influenza virus activity. *Proc Natl Acad Sci USA*. 2014;111:16820-5. (Surface Plasmon Resonance)
24. Damschroder MM, Widjaja L, Gill PS, Krasnoperov V, Jiang W, Dall'Acqua WF, et al. Framework shuffling of antibodies to reduce immunogenicity and manipulate functional and biophysical properties. *Mol Immunol*. 2007;44:3049-60. (Purification)
25. Su K, Li X, Ederberg JC, Wu J, Ferguson P, Kimberly RP. A promoter haplotype of the immunoreceptor tyrosine-based inhibitory motif-bearing FcγRIIb alters receptor expression and associates with autoimmunity. II. Differential binding of GATA4 and Yin-Yang1 transcription factors and correlated receptor expression and function. *J Immunol*. 2004;172:7192-9. (Stim)

Texas Red® is a registered trademark of Molecular Probes, Inc.

Alexa Fluor® 488, 647, and 555 are provided under an Intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.

TB2060

23-Aug-17

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA **Mailing Address:** P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • Fax: 205.945.8768

Email: info@southernbiotech.com • Website: www.southernbiotech.com