


## Goat Anti-Rabbit IgG(H+L), Mouse/Human ads

Cat. No.	Format	Size
4050-01	Purified (UNLB)	1.0 mg
4050-02	Fluorescein (FITC)	1.0 mg
4050-03	Rhodamine (TRITC)	1.0 mg
4050-04	Alkaline Phosphatase (AP)	1.0 mL
4050-05	Horseradish Peroxidase (HRP)	1.0 mL
4050-07	Texas Red <sup>®</sup> (TXRD)	1.0 mg
4050-08	Biotin (BIOT)	1.0 mg
4050-09	R-phycoerythrin (PE)	0.5 mg
4050-09S	R-phycoerythrin (PE)	0.25 mg
4050-11L	Allophycocyanin (APC)	0.5 mg
4050-11S	Allophycocyanin (APC)	0.25 mg
4050-13	Spectral Red <sup>®</sup> (SPRD)	0.25 mg
4050-30	Alexa Fluor <sup>®</sup> 488 (AF488)	1.0 mg
4050-31	Alexa Fluor <sup>®</sup> 647 (AF647)	1.0 mg
4050-32	Alexa Fluor <sup>®</sup> 555 (AF555)	1.0 mg



### Description

<b>Specificity</b>	Reacts with the heavy and light chains of rabbit IgG and the light chains of rabbit IgM
<b>Source</b>	Pooled antisera from goats hyperimmunized with rabbit IgG
<b>Cross Adsorption</b>	Mouse and human immunoglobulins and pooled sera; may react with immunoglobulins from other species
<b>Purification</b>	Affinity chromatography on pooled rabbit IgG covalently linked to agarose

### Applications

Quality tested applications include –

ELISA<sup>1-3</sup>  
FLISA  
FC<sup>4-7</sup>

Other referenced applications include –

IHC-FS<sup>8-10</sup>  
IHC-PS<sup>6,11-13</sup>  
IHC-WM<sup>14,15</sup>  
ICC<sup>16-19</sup>  
WB<sup>20-28</sup>  
Sep<sup>29</sup>

### Working Dilutions

<b>ELISA</b>	AP conjugate	1:2,000 – 1:4,000
	HRP conjugate	1:4,000 – 1:8,000
	BIOT conjugate	1:5,000 – 1:20,000
<b>FLISA</b>	FITC, TRITC, TXRD, AF488, and AF555 conjugates	1:100 – 1:400
	PE, APC, and AF647 conjugates	≤ 1 µg/mL
<b>Flow Cytometry</b>	FITC, BIOT, and AF488 conjugates	≤ 1 µg/10 <sup>6</sup> cells
	PE, APC, SPRD, and AF647 conjugates	≤ 0.1 µg/10 <sup>6</sup> cells
	For flow cytometry, the suggested use of these reagents is in a final volume of 100 µL	
<b>Other Applications</b>	Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.	

**For Research Use Only. Not for Diagnostic or Therapeutic Use.**

## Handling and Storage

- The purified (UNLB) antibody is supplied as 1.0 mg purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. No preservatives or amine-containing buffer salts added. Store at 2-8°C.
- The fluorescein (FITC), rhodamine (TRITC), Texas Red® (TXRD), Alexa Fluor® 488 (AF488), Alexa Fluor® 647 (AF647), and Alexa Fluor® 555 (AF555) conjugates are supplied as 1.0 mg in 1.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The alkaline phosphatase (AP) conjugate is supplied as 1.0 mL of stock solution in 50 mM Tris/1 mM MgCl<sub>2</sub>/50% glycerol, pH 8.0, containing NaN<sub>3</sub> as preservative. Store at 2-8°C or long-term at -20°C.
- The horseradish peroxidase (HRP) conjugate is supplied as 1.0 mL of stock solution in 50% glycerol/50% PBS, pH 7.4. No preservative added. Store at 2-8°C or long-term at -20°C.
- The biotin (BIOT) conjugate is supplied as 1.0 mg in 2.0 mL of PBS/NaN<sub>3</sub>. Store at 2-8°C.
- The R-phycoerythrin (PE) and allophycocyanin (APC) conjugates are supplied as 0.5 mg in 1.0 mL or 0.25 mg in 0.5 mL of PBS/NaN<sub>3</sub> and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- The Spectral Red® (SPRD) conjugate is supplied as 0.25 mg in 1.0 mL of PBS/NaN<sub>3</sub> and a stabilizing agent. Store at 2-8°C. **Do not freeze!**
- Protect fluorochrome-conjugated forms from light. Reagents are stable for the period shown on the label if stored as directed.

## Warning

Some reagents contain sodium azide. Please refer to product specific (M)SDS.

## References

1. Maroney AC, Finn JP, Connors TJ, Durkin JT, Angeles T, Gessner G, et al. Cep-1347 (KT7515), a semisynthetic inhibitor of the mixed lineage kinase family. *J Biol Chem.* 2001;276:25302-8. (ELISA)
2. Zhang X, He M, Cheng L, Chen Y, Zhou L, Zeng H, et al. Elevated heat shock protein 60 levels are associated with higher risk of coronary heart disease in Chinese. *Circulation.* 2008;118:2687-93. (ELISA)
3. Jin K, Wang S, Zhang C, Xiao Y, Lu S, Huang Z. Protective antibody responses against *Clostridium difficile* elicited by a DNA vaccine expressing the enzymatic domain of toxin B. *Hum Vaccin Immunother.* 2013;9:63-73. (ELISA)
4. Burger PE, Gupta R, Xiong X, Ontiveros CS, Salm SN, Moscatelli D, et al. High aldehyde dehydrogenase activity: a novel functional marker of murine prostate stem/progenitor cells. *Stem Cells.* 2009;27:2220-8. (FC)
5. Spurlock CF 3<sup>rd</sup>, Tossberg JT, Fuchs HA, Olsen NJ, Aune TM. Methotrexate increases expression of cell cycle checkpoint genes via JNK activation. *Arthritis Rheum.* 2012;64:1780-9. (FC)
6. Aziz M, Yang W, Matsuo S, Sharma A, Zhou M, Wang P. Upregulation of GRAIL is associated with impaired CD4 T cell proliferation in sepsis. *J Immunol.* 2014;192:2305-14. (FC, IHC-PS)
7. Boonstra MC, Tolner B, Schaafsma BE, Boogerd LS, Prevoo HA, Bhavsar G, et al. Preclinical evaluation of a novel CEA-targeting near-infrared fluorescent tracer delineating colorectal and pancreatic tumors. *Int J Cancer.* 2015;137:1910-20. (FC)
8. Borrego L, Maynard B, Peterson EA, George T, Iglesias L, Peters MS, et al. Deposition of eosinophil granule proteins precedes blister formation in bullous pemphigoid. Comparison with neutrophil and mast cell granule proteins. *Am J Pathol.* 1996;148:897-909. (IHC-FS)
9. Rønnov-Jessen L, Villadsen R, Edwards JC, Petersen OW. Differential expression of a chloride intracellular channel gene, CLIC4, in transforming growth factor- $\beta$  1-mediated conversion of fibroblasts to myofibroblasts. *Am J Pathol.* 2002;161:471-80. (IHC-FS)
10. Xia RH, Yosef N, Ubogu EE. Clinical, electrophysiological and pathologic correlations in a severe murine experimental autoimmune neuritis model of Guillain-Barré syndrome. *J Neuroimmunol.* 2010;219:54-63. (IHC-FS)
11. Penkowa M, Carrasco J, Giralt M, Molinero A, Hernández J, Campbell IL, et al. Altered central nervous system cytokine-growth factor expression profiles and angiogenesis in metallothionein-I-II deficient mice. *J Cereb Blood Flow Metab.* 2000;20:1174-89. (IHC-PS)
12. He X, Schoeb TR, Panoskaltis-Mortari A, Zinn KR, Kesterson RA, Zhang J, et al. Deficiency of P-selectin or P-selectin glycoprotein ligand-1 leads to accelerated development of glomerulonephritis and increased expression of CC chemokine ligand 2 in lupus-prone mice. *J Immunol.* 2006;177:8748-56. (IHC-PS)
13. Rossato E, Mkaddem SB, Kanamaru Y, Hurtado-Nedelec M, Hayem G, Descatoire V, et al. Reversal of arthritis by human monomeric IgA through the receptor-mediated SH2 domain-containing phosphatase 1 inhibitory pathway. *Arthritis Rheumatol.* 2015;67:1766-77. (IHC-PS)
14. Hens J, Gajda M, Scheuermann DW, Adriaensens D, Timmermans J. The longitudinal smooth muscle layer of the pig small intestine is innervated by both myenteric and submucous neurons. *Histochem Cell Biol.* 2002;117:481-92. (IHC-WM)
15. Monte E, Mouillesseaux K, Chen H, Kimball T, Ren S, Wang Y, et al. Systems proteomics of cardiac chromatin identifies nucleolin as a regulator of growth and cellular plasticity in cardiomyocytes. *Am J Physiol Heart Circ Physiol.* 2013;305:H1624-38. (IHC-WM)
16. Roberts MS, Woods AJ, Dale TC, van der Sluijs P, Norman JC. Protein kinase B/Akt acts via glycogen synthase kinase 3 to regulate recycling of  $\alpha$ v $\beta$ 3 and  $\alpha$ 5 $\beta$ 1 integrins. *Mol Cell Biol.* 2004;24:1505-15. (ICC)
17. Luijten MN, Basten SG, Claessens T, Vernooij M, Scott CL, Janssen R, et al. Birt-Hogg-Dubé syndrome is a novel ciliopathy. *Hum Mol Genet.* 2013;22:4383-97. (ICC)
18. Tarafdar S, Poe JA, Smithgall TE. The accessory factor Nef links HIV-1 to Tec/Btk kinases in an Src homology 3 domain-dependent manner. *J Biol Chem.* 2014;289:15718-28. (ICC)
19. Carey AJ, Tan CK, Mirza S, Irving-Rodgers H, Webb RI, Lam A, et al. Infection and cellular defense dynamics in a novel 17 $\beta$ -estradiol murine model of chronic human group B streptococcus genital tract colonization reveal a role for hemolysin in persistence and neutrophil accumulation. *J Immunol.* 2014;192:1718-31. (ICC)
20. Chen Z, Fu L, Herrero R, Schiffman M, Burk RD. Identification of a novel human papillomavirus (HPV97) related to HPV18 and HPV45. *Int J Cancer.* 2007;121:193-8. (WB)
21. Iuchi K, Hatano Y, Yagura T. Heterocyclic organobismuth(III) induces apoptosis of human promyelocytic leukemic cells through activation of caspases and mitochondrial perturbation. *Biochem Pharmacol.* 2008;76:974-86. (WB)
22. Fadnes B, Husebekk A, Svineng G, Rekdal Ø, Yanagishita M, Kolset SO, et al. The proteoglycan repertoire of lymphoid cells. *Glycoconj J.* 2012;29:513-23. (WB)
23. Tong J, Meyer JH, Furukawa Y, Boileau I, Chang L, Wilson AA, et al. Distribution of monoamine oxidase proteins in human brain: implications for brain imaging studies. *J Cereb Blood Flow Metab.* 2013;33:863-71. (WB)
24. Wu Y, Wang X, Guo H, Zhang B, Zhang X, Shi Z, et al. Synthesis and screening of 3-MA derivatives for autophagy inhibitors. *Autophagy.* 2013;9:595-603. (WB)
25. Au-Yeung BB, Melichar HJ, Ross JO, Cheng DA, Zikherman J, Shokat KM, et al. Quantitative and temporal requirements revealed for Zap70 catalytic activity during T cell development. *Nat Immunol.* 2014;15:687-94. (WB)
26. Goss JW, Kim S, Bledsoe H, Pollard TD. Characterization of the roles of Btl1p in fission yeast cytokinesis. *Mol Biol Cell.* 2014;25:1946-57. (WB)
27. Hogan MC, Bakeberg JL, Gainullin VG, Irazabal MV, Harmon AJ, Lieske JC, et al. Identification of biomarkers for PKD1 using urinary exosomes. *J Am Soc Nephrol.* 2015;26:1661-70. (WB)
28. Graf FE, Baker N, Munday JC, de Koning HP, Horn D, Mäser P. Chimerization at the AQP2-AQP3 locus is the genetic basis of melarsoprol-pentamidine cross-resistance in clinical *Trypanosoma brucei* gambiense isolates. *Int J Parasitol Drugs Drug Resist.* 2015;5:65-8. (WB)
29. Semba K, Namekata K, Guo X, Harada C, Harada T, Mitamura Y. Renin-angiotensin system regulates neurodegeneration in a mouse model of normal tension glaucoma. *Cell Death Differ.* 2014;5:e1333. (Sep)

Texas Red® is a registered trademark of Molecular Probes, Inc.

Spectral Red® is a registered trademark of Southern Biotechnology Associates, Inc.

Spectral Red® is a PE/CY5 tandem conjugate.

Cy5® is a registered trademark of GE Healthcare.

Alexa Fluor® 488, 647, and 555 are provided under an Intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes. Commercial Purposes means any activity by a party for consideration and may include, but is not limited to: (1) use of the product or its components in manufacturing; (2) use of the product or its components to provide a service, information, or data; (3) use of the product or its components for therapeutic, diagnostic or prophylactic purposes; or (4) resale of the product or its components, whether or not such product or its components are resold for use in research. For information on purchasing a license to this product for any other use, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.

TB4050

18-Jul-16

Corporate Offices: 160 Oxmoor Blvd • Birmingham, AL 35209 • USA Mailing Address: P.O. Box 26221 • Birmingham, AL 35260 • USA

Tel: 205.945.1774 • U.S. and Canada: 800.722.2255 • Fax: 205.945.8768

Email: [info@southernbiotech.com](mailto:info@southernbiotech.com) • Website: [www.southernbiotech.com](http://www.southernbiotech.com)