




Mouse Anti-Human IgE Fc

Cat. No.	Format	Size
9240-01	Purified (UNLB)	0.5 mg



ELISA plate was coated with purified human IgG, IgM, IgA, and IgE. Immunoglobulins were detected with serially diluted Mouse Anti-Human IgE Fc-UNLB (SB Cat. No. 9240-01) followed by Goat Anti-Mouse IgM, Human ads-HRP (SB Cat. No. 1020-05).

Overview

Clone	HP6061
Isotype	Mouse IgM κ
Immunogen	PS human IgE myeloma protein
Specificity	Human IgE Fc; Mr 188 kDa

Applications

ELISA – Quality tested ¹⁻⁶
 ELISPOT – Reported in literature ⁵
 Purification – Reported in literature ⁶
 Stim – Reported in literature ^{7,8}

Working Dilutions

ELISA	Purified (UNLB) antibody	2.5 – 10 µg/mL
Other Applications	Since applications vary, you should determine the optimum working dilution for the product that is appropriate for your specific need.	

For Research Use Only. Not for Diagnostic or Therapeutic Use.

Handling and Storage

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of borate buffered saline, pH 8.2. *No preservatives or amine-containing buffer salts added.* Store at 2-8°C.
- Reagents are stable for the period shown on the label if stored as directed.

References

1. Agea E, Russano A, Bistoni O, Mannucci R, Nicoletti I, Corazzi L, et al. Human CD1-restricted T cell recognition of lipids from pollens. *J Exp Med.* 2005;202:295-308. (ELISA)
2. Milovanovic M, Heine G, Zuberbier T, Worm M. Allergen extract-induced interleukin-10 in human memory B cells inhibits immunoglobulin E production. *Clin Exp Allergy.* 2009;39:671-8. (ELISA)
3. Heine G, Dahten A, Hilt K, Ernst D, Milovanovic M, Hartmann B, et al. Liver X receptors control IgE expression in B cells. *J Immunol.* 2009;182:5276-82. (ELISA)
4. Milovanovic M, Drozdenko G, Weise C, Babina M, Worm M. Interleukin-17A promotes IgE production in human B cells. *J Invest Dermatol.* 2010;130:2621-8. (ELISA)
5. Weise C, Hilt K, Milovanovic M, Ernst D, Rühl R, Worm M. Inhibition of IgE production by docosahexaenoic acid is mediated by direct interference with STAT6 and NFκB pathway in human B cells. *J Nutr Biochem.* 2011;22:269-75. (ELISA, ELISPOT)
6. Kleine-Tebbe J, Hamilton RG, Roebber M, Lichtenstein LM, MacDonald SM. Purification of immunoglobulin E (IgE) antibodies from sera with high IgE titers. *J Immunol Methods.* 1995;179:153-64. (Purification, ELISA)
7. Dölle S, Lehmann K, Schwarz D, Weckwert W, Scheler C, George E, et al. Allergenic activity of different tomato cultivars in tomato allergic subjects. *Clin Exp Allergy.* 2011;41:1643-52. (Stim)
8. Welter S, Dölle S, Lehmann K, Schwarz D, Weckwerth W, Worm M, et al. Pepino mosaic virus infection of tomato affects allergen expression, but not the allergenic potential of fruits. *PLoS One.* 2013;8(6):e65116. (Stim)