

Physalin L

Chemical Properties
CAS No.: 113146-74-0
Formula: C28H32O10
Molecular Weight: 528.55
Appearance: N/A
Storage: 0-4°C for short term (days to weeks), or -20°C for long term (mont

Biological Description

Description	Physalin L is a natural product, it shows a distinct fluorescence spot under UV 365 nm with good separation.		
Targets(IC ₅₀)	others: None		
In vitro	In this study, the inducible nitric oxide synthase (iNOS) assay was used to demonstrate that the dichloromethane extract of Physalis alkekengi var. franchetii (DCEP) possesses anti-inflammatory activity that might be attributed to the modification of key cysteine residues in IKK β by the MRAs in DCEP. To isolate these MRAs, glutathione (GSH) was employed, and a simple ultra-performance liquid chromatography/tandem mass spectrometry (UPLC-MS/MS) screening method was developed to investigate the GSH conjugates with potential MRAs. Five physalins, including one new compound isophysalin A (2), together with four known steroidal compounds, physalin A (1), physalin O (3), Physalin L (4) and physalin G (5), were isolated to evaluate the GSH conjugating abilities, and it was indicated that compounds 1, 2 and 3, which had a common $\alpha \pm ,\beta$ -unsaturated ketone moiety, exhibited conjugating abilities with GSH and also showed significant nitric oxide (NO) production inhibiting activities. The anti-inflammatory activities of compounds 1, 2 and 3 might be attributed to their targeting multiple cysteine residues on IKK β ; therefore, the alkylation of IKK β by compound 1 was further studied by micrOTOF-MS.		

Solubility Information

Solubility

< 1 mg/ml refers to the product slightly soluble or insoluble

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.892 mL	9.46 mL	18.92 mL
5 mM	0.378 mL	1.892 mL	3.784 mL
10 mM	0.189 mL	0.946 mL	1.892 mL
50 mM	0.038 mL	0.189 mL	0.378 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. The storage conditions and period of the stock solution: - 80 $^{\circ}$ C for 6 months; - 20 $^{\circ}$ C for 1 month. Please use it as soon as possible.

Reference

1. Physalins with anti-inflammatory activity are present in Physalis alkekengi var. franchetii and can function as Michael reaction acceptors. Steroids. 2012 Apr;77(5):441-7.

Inhibitors · Natural Compounds · Compound Libraries

This product is for Research Use Only \cdot Not for Human or Veterinary or Therapeutic Use.

Tel:781-999-4286 E-mail:info@targetmol.com Address:36 Washington Street,Wellesley Hills,MA 02481